

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
16 August 2001 (16.08.2001)

PCT

(10) International Publication Number
WO 01/58956 A2

(51) International Patent Classification⁷: **C07K 16/00**

(21) International Application Number: PCT/US01/04170

(22) International Filing Date: 9 February 2001 (09.02.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/181,608 10 February 2000 (10.02.2000) US

(63) Related by continuation (CON) or continuation-in-part (CIP) to earlier application:
US 60/181,608 (CON)
Filed on 10 February 2000 (10.02.2000)

(71) Applicant (for all designated States except US): **BASF AKTIENGESELLSCHAFT** [DE/DE]; Ludwigshafen, Rheinland-Pfalz (DE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **GHAYUR, Tariq** [PK/US]; 1014 Washington Street, Holliston, MA 01746 (US). **DIXON, Richard, W.** [US/US]; 6 Samuel Drive, North Grafton, MA 01546 (US). **ROGUSKA, Mike** [US/US]; 16 Hilldale Road, Ashland, MA 01721 (US). **WHITE, Michael** [US/US]; 30 Angelica Drive, Framingham, MA 01701 (US). **LABKOVSKY, Boris** [US/US]; 107A-5 Broadmeadow Road, Marlborough, MA 01752 (US). **SALFELD, Jochen** [DE/US]; 177 Old Westboro Road, North Grafton, MA 01536 (US). **DUNCAN, Alexander, Robert** [GB/GB]; 8 Garden Fields,

Little Shelford, Cambridge CB2 3HH (GB). **BROCK-LEHURST, Simon, Mark** [GB/GB]; 38 Huntmill, Fulbourn, Cambridge CB1 5RH (GB). **MANKOVICH, John** [US/US]; 416 Lowell Street, Andover, MA 01810 (US). **SHORROCK, Celia, Patricia** [GB/US]; 17 Stanley Road, Cambridge CB5 8LF (GB). **THOMPSON, Julia, Elizabeth** [GB/GB]; Seeby's Cottage, 6 High Street, Whittlesford, Cambridge CB2 4LT (GB). **LENNARD, Simon, Nicholas** [GB/GB]; 1 Bakers Lane, Linton, Cambridgeshire CB1 6NF (GB).

(74) Agents: **DECONTI, Giulio, A., JR.** et al.; Lahive & Cockfield, LLP, 28 State Street, Boston, MA 02109 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: ANTIBODIES THAT BIND HUMAN INTERLEUKIN-18 AND METHODS OF MAKING AND USING

(57) Abstract: Antibodies that bind human interleukin-18 (hIL-18) are provided, in particular antibodies that bind epitope(s) of human IL-18. The antibodies can be, for example, entirely human antibodies, recombinant antibodies, or monoclonal antibodies. Preferred antibodies have high affinity for hIL-18 and neutralize hIL-18 activity *in vitro* and *in vivo*. An antibody of the invention can be a full-length antibody or an antigen-binding portion thereof. Method of making and method of using the antibodies of the invention are also provided. The antibodies, or antibody portions, of the invention are useful for detecting hIL-18 and for inhibiting hIL-18 activity, e.g., in a human subject suffering from a disorder in which hIL-18 activity is detrimental.

BEST AVAILABLE COPY

ANTIBODIES THAT BIND HUMAN INTERLEUKIN-18 AND METHODS OF MAKING AND USING

Related Applications

5 This application is a non-provisional application claiming priority to U.S. provisional application Serial No. 60/181,608, filed February 10, 2000, entitled, "Antibodies that Bind Human Interleukin-18 and Methods of Making and Using" the contents of which are hereby incorporated by reference. In addition, the contents of all cited references, including literature references, issued patents, and published patent
10 applications, as cited throughout this application are hereby expressly incorporated by reference.

Background of the Invention

Interleukin-18 (IL-18) was originally described in 1989 as interferon-gamma
15 inducing factor (IGIF) and is a pro-inflammatory cytokine with various functions in addition to an ability to induce interferon gamma. These biological properties include activation of NF- κ b, Fas ligand expression, the induction of both CC and CXC chemokines, and increased production of competent human immunodeficiency virus.

Due to the ability of IL-18 to induce interferon gamma production in T cells and
20 macrophages, it plays an important role in Th1-type immune responses and participates in both innate and acquired immunity. IL-18 is related to the IL-1 family in terms of both structure and function. For reviews of IL-18 structure, function and biological activity, see for example Dinarello, C. et al. (1998) *J. Leukoc. Biol.* 63:658-654; Dinarello, C.A. (1999) *Methods* 19:121-132; and Dinarello, C.A. (1999) *J. Allergy Clin.*
25 *Immunol.* 103:11-24.

It would be desirable to use to modulate IL-18 in a variety of human immune responses. In particular, antibodies that bind to and neutralize IL-18 are particularly desirable. Moreover, murine IL-18 antibodies are limited for their use *in vivo* due to problems associated with administration of mouse antibodies to humans, such as short
30 serum half life, an inability to trigger certain human effector functions and elicitation of

- 2 -

an unwanted immune response against the mouse antibody in a human (the "human anti-mouse antibody" (HAMA) reaction).

In general, attempts to overcome the problems associated with use of fully-murine antibodies in humans, have involved genetically engineering the antibodies to be more "human-like." For example, chimeric antibodies, in which the variable regions of the antibody chains are murine-derived and the constant regions of the antibody chains are human-derived, have been prepared (Junghans, *et al.* (1990) *Cancer Res.* 50:1495-1502; Brown *et al.* (1991) *Proc. Natl. Acad. Sci.* 88:2663-2667; Kettleborough *et al.* (1991) *Protein Engineering.* 4:773-783). However, because these chimeric and humanized antibodies still retain some murine sequences, they still may elicit an unwanted immune reaction, the human anti-chimeric antibody (HAMA) reaction, especially when administered for prolonged periods.

A preferred IL-18 inhibitory agent to murine antibodies or derivatives thereof (*e.g.*, chimeric or humanized antibodies) would be an entirely human anti-IL-18 antibody, since such an agent should not elicit the HAMA reaction, even if used for prolonged periods. However, such antibodies have not been described in the art and, therefore are still needed.

Summary of the Invention

This invention pertains to compounds, such as antibodies, that bind human IL-18, as well as methods of making and using such compounds or antibodies.

In one aspect, the invention pertains to a compound capable of binding a human IL-18 amino acid sequence, or portion thereof, where the amino acid comprises an N- or C-terminal portion of human IL-18 such as provided in SEQ ID NO: 70 or SEQ ID NO: 71. In one embodiment, the compound is a small molecule, peptide, polypeptide, antibody, or antibody fragment, such as a fully human antibody or fragment.

In another aspect, the invention pertains to a human monoclonal antibody, or antigen-binding portion thereof, capable of binding to human IL-18. In other embodiments, the antibody or fragment thereof, dissociates from human IL-18, as determined by plasmon resonance, with a koff rate constant of 0.1s⁻¹ or less, 1 x 10E-2 s⁻¹ or less, 1 x 10E-3 s⁻¹ or less, 1 x 10E-4 s⁻¹ or less, 1 x 10E-5 s⁻¹ or less, 1 x 10E-6

- 3 -

s-1 or less, or inhibits human IL-18 activity with an IC₅₀ of 1 x 10E-6 or less, 1 x 10E-7 or less, 1 x 10E-8 or less, 1 x 10E-9, 1 x 10E-10 or less, or 1 x 10E-11 or less.

In another aspect, the invention pertains to an isolated antibody, or an antigen-binding portion thereof, that binds an epitope of human IL-18 comprising amino acids
5 PLFEDMTDSDCRDNA (SEQ ID NO: 1), VIRNLNDQVLFIDQ (SEQ ID NO: 33), or a portion of either. Preferably, the antibody is a neutralizing antibody. Preferably, the antibody is a human antibody. In various embodiments, the antibody is a recombinant antibody (*e.g.*, a single-chain antibody (scFv)), or a monoclonal antibody.

In other embodiments, the isolated antibody, or antigen-binding portion thereof,
10 binds to an epitope of human IL-18, or a portion of either, where the antibody, or antigen-binding portion thereof, dissociates from human IL-18 with a k_{off} rate constant of 0.1s⁻¹ or less, as determined by surface plasmon resonance, or which inhibits human IL-18 activity with an IC₅₀ of 1 x 10⁻⁶M or less. Alternatively, the antibody, or an antigen-binding portion thereof, may dissociate from human IL-18 with a k_{off} rate
15 constant of 1 x 10⁻²s⁻¹ or less, as determined by surface plasmon resonance, or may inhibit human IL-18 activity with an IC₅₀ of 1 x 10⁻⁷M or less. Alternatively, the antibody, or an antigen-binding portion thereof, may dissociate from human IL-18 with a k_{off} rate constant of 1 x 10⁻³s⁻¹ or less, as determined by surface plasmon resonance, or may inhibit human IL-18 activity with an IC₅₀ of 1 x 10⁻⁸M or less. Alternatively, the
20 antibody, or an antigen-binding portion thereof, may dissociate from human IL-18 with a k_{off} rate constant of 1 x 10⁻⁴s⁻¹ or less, as determined by surface plasmon resonance, or may inhibit human IL-18 activity with an IC₅₀ of 1 x 10⁻⁹M or less. Alternatively, the antibody, or an antigen-binding portion thereof, may dissociate from human IL-18 with a k_{off} rate constant of 1 x 10⁻⁵s⁻¹ or less, as determined by surface plasmon resonance, or
25 may inhibit human IL-18 activity with an IC₅₀ of 1 x 10⁻¹⁰M or less. Alternatively, the antibody, or an antigen-binding portion thereof, may dissociate from human IL-18 with a k_{off} rate constant of 1 x 10⁻⁶s⁻¹ or less, as determined by surface plasmon resonance, or may inhibit human IL-18 activity with an IC₅₀ of 1 x 10⁻¹¹M or less.

Another aspect of the invention pertains to an isolated human antibody, or an
30 antigen-binding portion thereof, containing at least one variable region CDR domain capable of binding an epitope of human IL-18. In related embodiments, the isolated

antibody, or an antigen-binding portion thereof, has a variable region containing a heavy and/or light chain CDR1 domain, CDR2 domain, or CDR3 domain as set forth in Table 6 or 9 which can have, *e.g.*, one or more amino acid substitutions or insertions at or adjacent to any of the Kabat positions indicated in Tables 7-8 and 10-11. In a preferred
5 embodiment, the isolated antibody, or an antigen-binding portion thereof, contains a light chain variable region (LCVR) containing the amino acid sequence of SEQ ID NO: 29 and a heavy chain variable region (HCVR) containing the amino acid sequence of SEQ ID NO: 26. In another preferred embodiment, the isolated antibody, or an antigen-binding portion thereof, contains a light chain variable region (LCVR) having the amino
10 acid sequence of SEQ ID NO: 29 and a heavy chain variable region (HCVR) having the amino acid sequence of SEQ ID NO: 27.

Another aspect of the invention pertains to pharmaceutical compositions comprising an antibody, or antigen-binding portion thereof, of the invention and a pharmaceutically acceptable carrier. In one embodiment, the pharmaceutical
15 composition further comprises at least one additional therapeutic agent for treating a disorder in which IL-18 activity is detrimental.

Another aspect of the invention pertains to methods of making an antibody that binds human interleukin-18 (IL-18). The invention provides a method comprising exposing an antibody repertoire to an antigen comprising an epitope of human IL-18
20 comprising amino acids PLFEDMTDSDCRDNA (SEQ ID NO: 1)), VIRNLNDQVLFIDQ (SEQ ID NO: 33), or a portion of either; and selecting from the antibody repertoire an antibody that binds the epitope of human IL-18 comprising amino acids PLFEDMTDSDCRDNA (SEQ ID NO: 1), VIRNLNDQVLFIDQ (SEQ ID NO: 33), or a portion of either.

25 In one embodiment, the antibody repertoire is an *in vivo* repertoire in an animal and the method comprises immunizing the animal with the antigen comprising the epitope of human IL-18 comprising amino acids PLFEDMTDSDCRDNA (SEQ ID NO: 1), VIRNLNDQVLFIDQ (SEQ ID NO: 33), the N- or C-terminal portion of human IL-18 (SEQ ID NOS: 70-71), or a portion of any of these epitopes. In another embodiment,
30 the antibody repertoire is a recombinant antibody library and the method comprises screening the library with an antigen containing the epitope of human IL-18 having the

- 5 -

amino acids PLFEDMTDSDCRDNA (SEQ ID NO: 1), VIRNLNDQVLFDQ (SEQ ID NO: 33), peptides represented by SEQ ID NOS: 31-32 and 34-60, or a portion of any of the foregoing. Preferably, the library is a human antibody library.

5 In another aspect, the invention provides an isolated nucleic acid encoding an antibody of any of the above aspects, *e.g.*, a heavy and/or light chain variable region, or portion thereof. In related embodiments, the isolated nucleic acid encoding the anti-IL-18 antibody, or portion thereof, is in a recombinant expression vector, *e.g.*, for expression in a host cell.

10 Thus, in another aspect, the invention pertains to a method of using the foregoing host cell into which the recombinant expression vector has been introduced, for synthesizing an antibody that binds human IL-18, by culturing the host cell in a culture medium until an antibody that binds human IL-18 is synthesized by the cell.

Another aspect of the invention pertains to a method for inhibiting human IL-18 activity comprising contacting human IL-18 with the antibody, or antigen-binding
15 portion thereof, of the invention such that human IL-18 activity is inhibited.

Yet another aspect of the invention pertains to a method for inhibiting human IL-18 activity in a human subject suffering from a disorder in which IL-18 activity is detrimental, comprising administering to the human subject the antibody, or antigen-binding portion thereof, of the invention such that human IL-18 activity in the human
20 subject is inhibited. In one embodiment, the anti-IL-18 antibody may be administered, *e.g.*, before, concurrent, or after, an additional agent such as an anti-IL-12 antibody or antigen binding fragment thereof, methotrexate, anti-TNF antibody or antigen binding fragment thereof, corticosteroids, cyclosporin, rapamycin, FK506, or a non-steroidal anti-inflammatory agent.

25

Brief Description of the Drawings

Figure 1 shows the structural model of IL-18 (center) as compared to IL-1 β (left) and IL1RA (right).

Figure 2 shows a structural model of IL-18 complexed with the IL-18 receptor, wherein the peptide epitope comprising amino acids PLFEDMTDSDCRDNA (SEQ ID NO: 1) of IL-18 is indicated in dark gray. This peptide epitope is bound by the anti-IL-18 antibody 2E1.

5

Figure 3 shows a structural model of IL-18 complexed with the IL-18 receptor, wherein the peptide epitope comprising amino acids YFGKLESKLSVIRN (SEQ ID NO: 33) of IL-18 is indicated in dark gray. This peptide epitope is bound by the anti-IL-18 antibody LT28.

10

Figure 4 shows a structural model of full length IL-18 complexed with the IL-18 receptor. The spherical light and dark gray epitopes represent the N and C terminal contact epitopes of IL-18 (respectively, SEQ ID NOS: 70 and 71).

15

Figure 5 shows the potency of three different anti-IL-18 antibodies in neutralizing the biologic effects of IL-18 as a function of inhibition of IFN- γ induction in KG1 cells. The IC₅₀ values for the antibodies 125H (boxes) and the 2E1 antibody as an IgG antibody (circles) or as a single chain antibody (triangles) are, respectively, 2.1E-10, 9.0E-10, and 3.3E-9.

20

Detailed Description of the Invention

This invention pertains to the selection of peptide epitopes that are capable of generating neutralizing antibodies to IL-18 mediated signal transduction, the preparation of antibodies to these epitopes and the use of such antibodies, including use to treat disorders involving IL-18. The strategy of selecting epitopes entails construction of an homology model of the IL-18 protein and its corresponding receptor. A combination of visual inspection and computational evaluation is then used to select representative peptide segments for synthesis and antibody generation. Amino acid sequences shown herein use the standard one-letter abbreviation code.

30

Selection of IL-18 Epitopes

- The program Modeler (Sali, A. et al., *Evaluation of comparative protein modeling by MODELLER*. Proteins: Struct., Funct., Genet. (1995), 23(3), pp. 318-26. CODEN: PSFGEY; ISSN: 0887-3585) was used to generate homology models for both
- 5 IL-18 and the IL-18 receptor. The X-ray crystal structures of IL-1 β (Priestle, J., et al. *The three-dimensional structure of human interleukin-1.beta. refined to 2.0 .ANG. resolution*. Prog. Clin. Biol. Res. (1990), 349 (Cytokines Lipocortins Inflammation Differ.), pp. 297-307) and IL-1RA (Schreuder, H. et al., *Refined crystal structure of the interleukin-1 receptor antagonist: presence of a disulfide link and a cis-proline*. Eur. J.
- 10 Biochem. (1995), 227(3), pp. 838-47) are available and were used as reference coordinates for the model construction of IL-18. The IL-1 receptor structure (Vigers, G., et al. *Crystal structure of the type-I interleukin-1-receptor complexed with interleukin-1.beta.* Nature (London) (1997), 386(6621), pp. 190-194) was used to model the IL-18 receptor.
- 15 The structural model building for IL-18 and the IL-18 receptor is described further below.

IL-18 Model Building

- The overall sequence homology with these two proteins (*i.e.*, IL-1 β and IL-18) is
- 20 low, however, there is compelling evidence that IL-18 is a member of the IL-1 family (see Dinarello, C.A. *IL-18: a TH1-inducing, proinflammatory cytokine and new member of the IL-1 family*. J. Allergy Clin. Immunol. (1999), 103(1, Pt. 1), pp. 11-24) and that the overall protein fold is very similar. Like IL-1 β , IL-18 is initially secreted in a pro form. Both pro-IL-1 β and pro-IL-18 are activated by IL-1 β -converting enzyme
- 25 (ICE) (Fantuzzi, G. and Dinarello, C.A. *Interleukin-18 and interleukin-1 β : two cytokine substrates for ICE (caspase-1)*. J. Clin. Immunol. (1999), 19(1), pp. 1-11). It is also known that the IL-1 receptor and the IL-18 receptor are similar (Dinarello, C.A. et al. *Overview of interleukin-18: more than an interferon- γ inducing factor*. J. Leukocyte Biol. (1998), 63(6), 658-664). IL-1 β is capable of binding to the IL-18 receptor. As a
- 30 final argument, IL-1 β and IL-1RA display an identical fold, even though the overall

sequence homology between these two proteins is on a par with the sequence homology with IL-18. The sequence alignment between the three proteins (*i.e.*, IL-18, IL-1 β and IL1-RA) was constructed manually with the program InsightII. This alignment can be seen in Table 1:

Table 1: Sequence alignment for IL-18 relative to IL-1 β and IL-1RA

		24	
5	IL-18:	YFGK-LESKLS-VIRNLNDQVLFIDQGNRPLFE--DMT-DSDCRD--NAP	
	IL-1 β :	AP-VRS-LNCTLRDSQQKSLVMS-G--P-YELKALHLQGQ--D--MEQ	
	IL-1RA:	SSKMQA-FR--IWDVNQKTFYLR-N--N--QLVAGYLQGP--NVNLEE	
		80	
10	IL-18:	RTIFIISMY-KDSQPRG-MAVTISVKCEKISTLSC----ENK-IISFKEM	
	IL-1 β :	QVVFSMS-FVQGEESNDKIPVALGLK-EKNLYLSCVLK-DDKPTLQLESV	
	IL-1RA:	KI--DV---VP-IEPH---ALFLGIH-GGKMCLSCV-KSGDETRLQLEAV	
		123	
15	IL-18:	NPPDNI-KDTKSDIIF-FQRSVPGHDNKMQFESSYEGYFLACE-KERDL	
	IL-1 β :	DPKNYP-KK-KMEKRFFVFNK-I-EINNKLFEFESAQFPNWIYSTS-QAENM	
	IL-1RA:	NITDLSENR-KQDKRFAFIR-S-DSGPTTSFESAACPGWFLCTAMEADQ-	
		170	
20	IL-18:	FKLILKKED-ELGDRSIM-FTVQNE	(SEQ ID NO: 4)
	IL-1 β :	-PVFL--GG-TKGGQDITDFTMQFVSS	(SEQ ID NO: 5)
	IL-1RA:	-PVSL--TNMPDEGVMVTKFYFQED	(SEQ ID NO: 6)

The sequence homology between these sequences is listed in Table 2. The upper triangle is percent strict sequence identity and the lower triangle is percent conservative sequence homology. Only the portions of the total sequences reported in Table 1 are considered in Table 2. As was mentioned above, the overall homology is low but consistent across the family.

Table 2: Sequence homology between IL-1 family members

Molecule	IL-18	IL-1β	IL-1RA	IL-1 Rec	IL-18 Rec
IL-18	-	20.0	21.8	-	-
IL-1b	53.5	-	27.5	-	-
IL-1RA	50.6	54.4	-	-	-
IL-1 Rec	-	-	-	-	26.1
IL-18 Rec	-	-	-	50.5	-

The resulting IL-18 structure is pictured in Figure 1 along with IL-1 β and IL-1RA. The overall quality as assessed by the program What_Check (Hoof, R.W. et al., *Errors in Protein Structures*. Nature (1996) 381, pp. 272) is reasonable, but a bit low (see Table 3 below).

Table 3: Structural Z-scores from What_Check (Positive is better than average)

	IL-1 β	IL-1RA	IL-1 Rec	IL-18 (model)	IL-18 Rec (model)
Packing Quality	-1.6	-2.3	-3.6	-5.5	-5.7
Ramachandran Plot	-2.0	-1.3	-2.9	-3.3	-2.3
Rotamer Normality	-1.6	-0.8	-1.5	-0.6	-0.6
Backbone Conformation	-1.7	+0.5	-1.6	-5.6	-2.7

5

However, the assessment of the reference structures by What_Check is also low, suggesting that this protein fold is poorly represented in the database of reference structures. Undoubtedly though, the low sequence homology contributed to a less than perfect final structure in spite of our confidence in the overall protein fold. However, for the purpose of choosing epitopes for antibody generation, this structure is considered to be sufficient.

10

IL-18 Receptor Model Building

The structure of the IL-18 receptor was also generated using the program Modeler. The reference coordinates were from the IL-1 receptor. As in the case of the cytokines associated with these receptors, the overall sequence identity is low, but sufficient to generate an alignment. The sequence homology figures are included in Table 2 above. The alignment was generated manually using the program InsightII and is presented in Table 4.

20

Table 4. Sequence alignment for the IL-18 receptor relative to the IL-1 Receptor

		22
	IL-18 Rec:	CTSRPHITVVEGEPFYLKHCSCSLAHEIETTTKSWYKSSGSQEHVI:LNPR
5	IL-1 Rec :	CKEREEKIILVSSANEIDVRPCPLNPNEHKGTITWYKDD-SKTPVSTEQA
		72
	IL-18 Rec:	SSSRIALHDCVLEFWPVELNDTGSYFFQMKNYTQKWKLNVIRRNKHS----
	IL-1 Rec :	S--RIHQHKEKLWFVPAKVEDSGHYCVVRNSSYCLRIKISAKFVENEPN
		119
	IL-18 Rec:	-CFTERQVTSKIVEVKKFFQITCENSYYQTLVNST-----SLYKNCKKLLL
	IL-1 Rec :	LCYNAQAIFKQKLPVAGDGLVCPYMEFFKNENNELPKLQWYKDCKPLLL
		163
	IL-18 Rec:	EN-----NKNPTIKKNAEFEDQGYYSVCVHFLHHNGKLFNITKTFNITIVE
	IL-1 Rec :	DNIHFSGVKDRLIVMNVAEKHRGNYTCHASYTYLGKQYPITRVIEFITLE
		209
10	IL-18 Rec:	DRSNIVPVLLGPKLNHVAVELGKNVRLNCSALLNEEDVIYWMF-GEE-NG
	IL-1 Rec :	ENKPTRPVIVSPANETMEVDLGSQIQLICNVTGQLSDIAYWKWNGSVIDE
		257
	IL-18 Rec:	SDPNIHEE-KEMRIMTPEGKWHASKVLRIENIGESNLNVLYNCTVASTGG
	IL-1 Rec :	DDPVLGEDYYSVENPANKRRSTLITVLNISEIESRIFYKHPFTCFKANTHG
		306
	IL-18 Rec:	TDTKSFILVRKAD (SEQ ID NO: 7)
	IL-1 Rec :	IDAAYIQLIYPVT (SEQ ID NO: 8)

15

The overall quality of the structure of the IL-18 receptor as determined using the Modeler program is reasonable but again scores somewhat low according to What_Check (see Table 3 above). The confidence that can be placed on the overall fold comes primarily from the fact that IL-1 β binds to both the IL-1 and IL-18 receptor. The low sequence homology certainly contributes to the quality of the final structure, however, as in the case of the associated cytokines above, this current structure is considered to be sufficient. As an additional exercise, the IL-18 peptide epitope bound by LT28 (SEQ ID NO: 33) was modeled when complexed with the IL-18 receptor (Figure 3). As a final exercise, a model of the IL-18/IL-18 receptor complex was generated based on the IL-1 β /IL-1 receptor structure (Figure 4). This structure was generated by superimposing the cytokine structures and the receptor structures. No attempt was made to energy minimize the final structure.

Peptide Epitope Selection

The primary purpose of generating structural models was to be able to select suitable peptides based primarily on a visual score. Solvent exposed sections of the

proteins and portions of the proteins which were both hydrophilic and buried in the receptor/cytokine complex were considered. A final element considered was based on a selectivity criterion. The selected peptide epitope should be different in sequence from similar portions of other members of the family. Based on this criteria, a peptide from

5 IL-18 was selected. In addition, a comprehensive overlapping panel of peptides (SEQ ID NOS: 31-60) representative of full length human IL-18 (SEQ ID NO: 61) was also made and the sequence of all of these IL-18 related peptides is shown in Table 5, below.

Table 5. Selected Peptides Representative of IL-18

10

Peptide Sequence	SEQ ID NO:
PLFEDMTDSDCRDNA	(SEQ ID NO: 1)
CPLFEDMTDSDCRDNA	(SEQ ID NO: 2)
PLFEDMTDSDCR	(SEQ ID NO: 3)
YFGKLESKLSVIRN	(SEQ ID NO: 31)
ESKLSVIRNLNDQV	(SEQ ID NO: 32)
VIRNLNDQVLFIDQ (LT28 binding epitope)	(SEQ ID NO: 33)
NDQVLFIDQGNRPL	(SEQ ID NO: 34)
FIDQGNRPLFEDMT	(SEQ ID NO: 35)
NRPLFEDMTDSDCR (2E1 binding epitope)	(SEQ ID NO: 36)
EDMTDSDCRDNAPR	(SEQ ID NO: 37)
SDCRDNAPRTIFII	(SEQ ID NO: 38)
NAPRTIFIISMYKD	(SEQ ID NO: 39)
IFIISMYKDSQPRG	(SEQ ID NO: 40)
MYKDSQPRGMAVTI	(SEQ ID NO: 41)
QPRGMAVTISVKCE	(SEQ ID NO: 42)
AVTISVKCEKISTL	(SEQ ID NO: 43)
VKCEKISTLSCENK	(SEQ ID NO: 44)
ISTLSCENKIISFK	(SEQ ID NO: 45)
CENKIISFKEMNPP	(SEQ ID NO: 46)

- 14 -

ISFKEMNPPDNIKD	(SEQ ID NO: 47)
MNPPDNIKDTKSDI	(SEQ ID NO: 48)
NIKDTKSDIIFQQR	(SEQ ID NO: 49)
KSDIIFQQRSVPGH	(SEQ ID NO: 50)
FFQQRVPGHDKMQ	(SEQ ID NO: 51)
VPGHDKMQFESSS	(SEQ ID NO: 52)
NKMQFESSSYEGYF	(SEQ ID NO: 53)
ESSSYEGYFLACEK	(SEQ ID NO: 54)
EGYFLACEKERDLF	(SEQ ID NO: 55)
ACEKERDLFKLILK	(SEQ ID NO: 56)
RDLFKLILKKEDEL	(SEQ ID NO: 57)
LILKKEDELGDRSI	(SEQ ID NO: 58)
EDELGDRSIMFTVQ	(SEQ ID NO: 59)
DRSIMFTVQNE	(SEQ ID NO: 60)
YFGKLESKLS VIRNLNDQVLFIDQGNRPLFEDMTD SDCRD NAPTIFIISMYKDSQPRGMAVTISVKCEK ISTLSCENKIISFKEMNPPDNIKDTKSDIIFQQR VPGHDKMQFESSSYEGYFLACEKERDLFKLILKK EDELGDRSIMFTVQNE	(SEQ ID NO: 61) LT28 and 2E1 epitopes indicated in bold

The N-terminal cysteine of the IL-18 peptide represented by SEQ ID NO: 2 is not part of the native IL-18 sequence, but was added as a conjugation site. Accordingly, within the native IL-18 amino acid sequence, the region corresponding to the selected epitope comprises amino acid residues having the amino acid sequence

5 PLFEDMTDSDCRDNA (SEQ ID NO: 1).

A schematic model of the IL-18 peptide (SEQ ID NO: 1) complexed with the IL-18 receptor is shown in Figure 2, with this peptide epitope indicated in dark gray.

Subsequent antigenicity calculations were performed on IL-18 peptide

10 sequences, with the result that this peptide scored particularly highly. This peptide was synthesized and used as an epitope to generate antibodies in a rabbit host. Molecular modeling data obtained using the IL-18 peptide PLFEDMTDSDCRDNA (SEQ ID NO: 1) or YFGKLESKLSVIRN (SEQ ID NO: 31) as compared to a cognate receptor, *i.e.*,

- 15 -

the IL-18 receptor, as depicted in Figures 2 and 3, provides an indication as to what residues a neutralizing antibody or compound may interact with.

An alternative method for peptide epitope selection can be accomplished in the absence of any molecular modeling by screening a panel of representative peptides using immunoselection. In one approach, overlapping peptides representative of the entire protein sequence can be used. In a more limited approach, only certain epitopes are represented in the panel of peptides. In a combined approach, molecular modeling can be used to identify epitopes likely to be important. The identified epitope(s) sequence can then be used to construct a panel of peptides (*e.g.*, overlapping peptides) that are representative of the identified epitope(s). Methods for manufacturing desired peptide sequences can be carried out using standard techniques known in the art.

Once the binding peptide or peptides (*e.g.*, panel of overlapping peptides) has been selected, an immunoscreen for a cognate receptor can be performed. Alternatively, an immunoscreen can be performed with a selected cognate receptor such that a peptide having a certain binding affinity can be identified. Any number of immunoscreens can be employed such that, either a desired receptor or desired peptide can be identified as a candidate binding molecule for further study. Such "bait" and "prey" techniques for analyzing protein-protein interactions, for identifying candidate binding molecules, and/or for scoring binding affinities are described in the art. One preferred technique utilizes phage display as described herein.

Anti-IL-18 Antibodies

The invention provides antibodies, as well as antibody portions thereof, that bind IL-18. Preferably, the antibodies, or portions thereof, are isolated antibodies.

Preferably, the antibodies, or portions thereof, are neutralizing antibodies.

The term "antibody", as used herein, is intended to refer to immunoglobulin molecules comprised of four polypeptide chains, two heavy (H) chains and two light (L) chains inter-connected by disulfide bonds. Each heavy chain is comprised of a heavy chain variable region (abbreviated herein as HCVR or VH) and a heavy chain constant region. The heavy chain constant region is comprised of three domains, CH1, CH2 and CH3. Each light chain is comprised of a light chain variable region (abbreviated herein

- 16 -

as LCVR or VL) and a light chain constant region. The light chain constant region is comprised of one domain, CL. The VH and VL regions can be further subdivided into regions of hypervariability, termed complementarity determining regions (CDR), interspersed with regions that are more conserved, termed framework regions (FR).

- 5 Each VH and VL is composed of three CDRs and four FRs, arranged from amino-terminus to carboxy-terminus in the following order: FR1, CDR1, FR2, CDR2, FR3, CDR3, FR4.

The term "antigen-binding portion" of an antibody (or simply "antibody portion"), as used herein, refers to one or more fragments of an antibody that retain the ability to specifically bind to an antigen (*e.g.*, hIL-18). It has been shown that the antigen-binding function of an antibody can be performed by fragments of a full-length antibody. Examples of binding fragments encompassed within the term "antigen-binding portion" of an antibody include (i) a Fab fragment, a monovalent fragment consisting of the VL, VH, CL and CH1 domains; (ii) a F(ab')₂ fragment, a bivalent fragment comprising two Fab fragments linked by a disulfide bridge at the hinge region; (iii) a Fd fragment consisting of the VH and CH1 domains; (iv) a Fv fragment consisting of the VL and VH domains of a single arm of an antibody, (v) a dAb fragment (Ward *et al.*, (1989) *Nature* 341:544-546), which consists of a VH domain; and (vi) an isolated complementarity determining region (CDR). Furthermore, although the two domains of the Fv fragment, VL and VH, are coded for by separate genes, they can be joined, using recombinant methods, by a synthetic linker that enables them to be made as a single protein chain in which the VL and VH regions pair to form monovalent molecules (known as single chain Fv (scFv); see *e.g.*, Bird *et al.* (1988) *Science* 242:423-426; and Huston *et al.* (1988) *Proc. Natl. Acad. Sci. USA* 85:5879-5883). Such single chain antibodies are also intended to be encompassed within the term "antigen-binding portion" of an antibody. Other forms of single chain antibodies, such as diabodies are also encompassed. Diabodies are bivalent, bispecific antibodies in which VH and VL domains are expressed on a single polypeptide chain, but using a linker that is too short to allow for pairing between the two domains on the same chain, thereby forcing the domains to pair with complementary domains of another chain and creating two antigen

10
15
20
25
30

binding sites (see *e.g.*, Holliger, P., *et al.* (1993) *Proc. Natl. Acad. Sci. USA* 90:6444-6448; Poljak, R.J., *et al.* (1994) *Structure* 2:1121-1123).

Still further, an antibody or antigen-binding portion thereof may be part of a larger immunoadhesion molecules, formed by covalent or noncovalent association of the antibody or antibody portion with one or more other proteins or peptides. Examples of such immunoadhesion molecules include use of the streptavidin core region to make a tetrameric scFv molecule (Kipriyanov, S.M., *et al.* (1995) *Human Antibodies and Hybridomas* 6:93-101) and use of a cysteine residue, a marker peptide and a C-terminal polyhistidine tag to make bivalent and biotinylated scFv molecules (Kipriyanov, S.M., *et al.* (1994) *Mol. Immunol.* 31:1047-1058). Antibody portions, such as Fab and F(ab')₂ fragments, can be prepared from whole antibodies using conventional techniques, such as papain or pepsin digestion, respectively, of whole antibodies. Moreover, antibodies, antibody portions and immunoadhesion molecules can be obtained using standard recombinant DNA techniques, as described herein.

An "isolated antibody", as used herein, is intended to refer to an antibody that is substantially free of other antibodies having different antigenic specificities (*e.g.*, an isolated antibody that specifically binds hIL-18 is substantially free of antibodies that specifically bind antigens other than hIL-18). An isolated antibody that specifically binds hIL-18 may, however, have cross-reactivity to other antigens, such as IL-18 molecules from other species. Moreover, an isolated antibody may be substantially free of other cellular material and/or chemicals. Further, an isolated antibody, *e.g.*, an isolated human antibody, can be a chimeric antibody wherein, *e.g.*, variable regions, CDR domains, or isotypes derived from a different human source are grafted to the parent human antibody.

A "compound" as used herein, refers to binding molecules such as antibodies, *e.g.*, polyclonal antibodies, monoclonal antibodies, binding fragments thereof (*e.g.*, Fab fragments), single chain antibodies (*e.g.*, scFv), peptides or peptide mimetics, as well as non-peptide based molecules, such as small molecules having ligand binding activity.

A "neutralizing antibody", as used herein (or an "antibody that neutralized hIL-18 activity"), is intended to refer to an antibody whose binding to hIL-18 results in inhibition of the biological activity of hIL-18. This inhibition of the biological activity

of hIL-18 can be assessed by measuring one or more indicators of hIL-18 biological activity. These indicators of hIL-18 biological activity can be assessed by one or more of several standard *in vitro* or *in vivo* assays known in the art.

The term "surface plasmon resonance", as used herein, refers to an optical
5 phenomenon that allows for the analysis of real-time biospecific interactions by detection of alterations in protein concentrations within a biosensor matrix, for example using the BIAcore system (Pharmacia Biosensor AB, Uppsala, Sweden and Piscataway, NJ). For further descriptions, see Jönsson, U., *et al.* (1993) *Ann. Biol. Clin.* 51:19-26; Jönsson, U., *et al.* (1991) *Biotechniques* 11:620-627; Johnsson, B., *et al.* (1995) *J. Mol.*
10 *Recognit.* 8:125-131; and Johnsson, B., *et al.* (1991) *Anal. Biochem.* 198:268-277.

The term " K_{off} ", as used herein, is intended to refer to the off rate constant for dissociation of an antibody from the antibody/antigen complex.

The term " K_d ", as used herein, is intended to refer to the dissociation constant of a particular antibody-antigen interaction.

15 In one aspect, the invention pertains to an isolated antibody, or an antigen-binding portion thereof, that binds an epitope of human IL-18 comprising amino acids PLFEDMTDSDCRDNA (SEQ ID NO: 1) or VIRNLNDQVLFIDQ (SEQ ID NO: 33), or a portion of any of these epitopes. Preferably, the antibody is a neutralizing antibody. Preferably, the antibody is a human antibody. In various embodiments, the antibody is a
20 recombinant antibody or a monoclonal antibody.

In other embodiments, the isolated antibody, or antigen-binding portion thereof, binds to an epitope of human IL-18 comprising amino acids PLFEDMTDSDCRDNA (SEQ ID NO: 1), wherein the antibody, or antigen-binding portion thereof, dissociates from human IL-18 with a k_{off} rate constant of $0.1s^{-1}$ or less, as determined by surface
25 plasmon resonance, or which inhibits human IL-18 activity with an IC_{50} of $1 \times 10^{-6}M$ or less. Alternatively, the antibody, or an antigen-binding portion thereof, may dissociate from human IL-18 with a k_{off} rate constant of $1 \times 10^{-2}s^{-1}$ or less, as determined by surface plasmon resonance, or may inhibit human IL-18 activity with an IC_{50} of $1 \times 10^{-7}M$ or less. Alternatively, the antibody, or an antigen-binding portion thereof, may dissociate
30 from human IL-18 with a k_{off} rate constant of $1 \times 10^{-3}s^{-1}$ or less, as determined by surface plasmon resonance, or may inhibit human IL-18 activity with an IC_{50} of $1 \times 10^{-8}M$ or

- 19 -

less. Alternatively, the antibody, or an antigen-binding portion thereof, may dissociate from human IL-18 with a k_{off} rate constant of $1 \times 10^{-4} \text{s}^{-1}$ or less, as determined by surface plasmon resonance, or may inhibit human IL-18 activity with an IC_{50} of $1 \times 10^{-9} \text{M}$ or less. Alternatively, the antibody, or an antigen-binding portion thereof, may dissociate from human IL-18 with a k_{off} rate constant of $1 \times 10^{-5} \text{s}^{-1}$ or less, as determined by surface plasmon resonance, or may inhibit human IL-18 activity with an IC_{50} of $1 \times 10^{-10} \text{M}$ or less. Alternatively, the antibody, or an antigen-binding portion thereof, may dissociate from human IL-18 with a k_{off} rate constant of $1 \times 10^{-6} \text{s}^{-1}$ or less, as determined by surface plasmon resonance, or may inhibit human IL-18 activity with an IC_{50} of $1 \times 10^{-11} \text{M}$ or less.

Affinity Maturation of Identified Anti-IL-18 Antibodies

The invention also provides for the further modification of an antibody identified as binding to an IL-18 epitope. The modification of the identified anti-IL-18 antibody is to improve binding and/or neutralization activity.

Therapeutic Compositions and Methods for Administering

The invention also provides pharmaceutical compositions comprising an antibody, or antigen-binding portion thereof, of the invention and a pharmaceutically acceptable carrier. In one embodiment, the pharmaceutical composition further comprises at least one additional therapeutic agent for treating a disorder in which IL-18 activity is detrimental.

The antibodies and antibody-portions of the invention can be incorporated into pharmaceutical compositions suitable for administration to a subject. Typically, the pharmaceutical composition comprises an antibody or antibody portion of the invention and a pharmaceutically acceptable carrier. As used herein, "pharmaceutically acceptable carrier" includes any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like that are physiologically compatible. Examples of pharmaceutically acceptable carriers include one or more of water, saline, phosphate buffered saline, dextrose, glycerol, ethanol and the like, as well as combinations thereof. In many cases, it will be preferable to include

- 20 -

isotonic agents, for example, sugars, polyalcohols such as mannitol, sorbitol, or sodium chloride in the composition. Pharmaceutically acceptable carriers may further comprise minor amounts of auxiliary substances such as wetting or emulsifying agents, preservatives or buffers, which enhance the shelf life or effectiveness of the antibody or
5 antibody portion.

The antibodies and antibody-portions of the invention can be incorporated into a pharmaceutical composition suitable for parenteral administration. Preferably, the antibody or antibody-portions will be prepared as an injectable solution containing 0.1-250 mg/ml antibody. The injectable solution can be composed of either a liquid or
10 lyophilized dosage form in a flint or amber vial, ampule or pre-filled syringe. The buffer can be L-histidine (1-50 mM), optimally 5-10 mM, at pH 5.0 to 7.0 (optimally pH 6.0). Other suitable buffers include but are not limited to, sodium succinate, sodium citrate, sodium phosphate or potassium phosphate. Sodium chloride can be used to modify the toxicity of the solution at a concentration of 0-300 mM (optimally 150 mM for a liquid
15 dosage form). Cryoprotectants can be included for a lyophilized dosage form, principally 0-10% sucrose (optimally 0.5-1.0%). Other suitable cryoprotectants include trehalose and lactose. Bulking agents can be included for a lyophilized dosage form, principally 1-10% mannitol (optimally 2-4%). Stabilizers can be used in both liquid and lyophilized dosage forms, principally 1-50 mM L-Methionine (optimally 5-10 mM).
20 Other suitable bulking agents include glycine, arginine, can be included as 0-0.05% polysorbate-80 (optimally 0.005-0.01%). Additional surfactants include but are not limited to polysorbate 20 and BRIJ surfactants.

The compositions of this invention may be in a variety of forms. These include, for example, liquid, semi-solid and solid dosage forms, such as liquid solutions (*e.g.*,
25 injectable and infusible solutions), dispersions or suspensions, tablets, pills, powders, liposomes and suppositories. The preferred form depends on the intended mode of administration and therapeutic application. Typical preferred compositions are in the form of injectable or infusible solutions, such as compositions similar to those used for passive immunization of humans with other antibodies. The preferred mode of
30 administration is parenteral (*e.g.*, intravenous, subcutaneous, intraperitoneal, intramuscular). In a preferred embodiment, the antibody is administered by intravenous

infusion or injection. In another preferred embodiment, the antibody is administered by intramuscular or subcutaneous injection.

Therapeutic compositions typically must be sterile and stable under the conditions of manufacture and storage. The composition can be formulated as a solution,
5 microemulsion, dispersion, liposome, or other ordered structure suitable to high drug concentration. Sterile injectable solutions can be prepared by incorporating the active compound (*i.e.*, antibody or antibody portion) in the required amount in an appropriate solvent with one or a combination of ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the active
10 compound into a sterile vehicle that contains a basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile, lyophilized powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and spray-drying that yields a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof. The
15 proper fluidity of a solution can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. Prolonged absorption of injectable compositions can be brought about by including in the composition an agent that delays absorption, for example, monostearate salts and gelatin.

20 The antibodies and antibody-portions of the present invention can be administered by a variety of methods known in the art, although for many therapeutic applications, the preferred route/mode of administration is subcutaneous injection, intravenous injection or infusion. As will be appreciated by the skilled artisan, the route and/or mode of administration will vary depending upon the desired results. In certain embodiments, the
25 active compound may be prepared with a carrier that will protect the compound against rapid release, such as a controlled release formulation, including implants, transdermal patches, and microencapsulated delivery systems. Biodegradable, biocompatible polymers can be used, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters, and polylactic acid. Many methods for the preparation of such
30 formulations are patented or generally known to those skilled in the art. See, *e.g.*,

Sustained and Controlled Release Drug Delivery Systems, J.R. Robinson, ed., Marcel Dekker, Inc., New York, 1978.

In certain embodiments, an antibody or antibody portion of the invention may be orally administered, for example, with an inert diluent or an assimilable edible carrier.

- 5 The compound (and other ingredients, if desired) may also be enclosed in a hard or soft shell gelatin capsule, compressed into tablets, or incorporated directly into the subject's diet. For oral therapeutic administration, the compounds may be incorporated with excipients and used in the form of ingestible tablets, buccal tablets, troches, capsules, elixirs, suspensions, syrups, wafers, and the like. To administer a compound of the
10 invention by other than parenteral administration, it may be necessary to coat the compound with, or co-administer the compound with, a material to prevent its inactivation.

- Supplementary active compounds can also be incorporated into the compositions. In certain embodiments, an antibody or antibody portion of the invention
15 is coformulated with and/or coadministered with one or more additional therapeutic agents that are useful for treating disorders in which IL-18 activity is detrimental. For example, an anti-hIL-18 antibody or antibody portion of the invention may be coformulated and/or coadministered with one or more additional antibodies that bind other targets (e.g., antibodies that bind other cytokines or that bind cell surface
20 molecules). Furthermore, one or more antibodies of the invention may be used in combination with two or more of the foregoing therapeutic agents. Such combination therapies may advantageously utilize lower dosages of the administered therapeutic agents, thus avoiding possible toxicities or complications associated with the various monotherapies.

25

Therapeutic Uses

- Interleukin 18 plays a critical role in the pathology associated with a variety of diseases involving immune and inflammatory elements. These diseases include, but are not limited to, rheumatoid arthritis, osteoarthritis, juvenile chronic arthritis, Lyme
30 arthritis, psoriatic arthritis, reactive arthritis, spondyloarthropathy, systemic lupus erythematosus, Crohn's disease, ulcerative colitis, inflammatory bowel disease, insulin

- dependent diabetes mellitus, thyroiditis, asthma, allergic diseases, psoriasis, dermatitis scleroderma, graft versus host disease, organ transplant rejection, acute or chronic immune disease associated with organ transplantation, sarcoidosis, atherosclerosis, disseminated intravascular coagulation, Kawasaki's disease, Grave's disease, nephrotic
- 5 syndrome, chronic fatigue syndrome, Wegener's granulomatosis, Henoch-Schoenlein purpura, microscopic vasculitis of the kidneys, chronic active hepatitis, uveitis, septic shock, toxic shock syndrome, sepsis syndrome, cachexia, infectious diseases, parasitic diseases, acquired immunodeficiency syndrome, acute transverse myelitis, Huntington's chorea, Parkinson's disease, Alzheimer's disease, stroke, primary biliary cirrhosis,
- 10 hemolytic anemia, malignancies, heart failure, myocardial infarction, Addison's disease, sporadic, polyglandular deficiency type I and polyglandular deficiency type II, Schmidt's syndrome, adult (acute) respiratory distress syndrome, alopecia, alopecia areata, seronegative arthropathy, arthropathy, Reiter's disease, psoriatic arthropathy, ulcerative colitic arthropathy, enteropathic synovitis, chlamydia, yersinia and salmonella
- 15 associated arthropathy, spondyloarthopathy, atheromatous disease/arteriosclerosis, atopic allergy, autoimmune bullous disease, pemphigus vulgaris, pemphigus foliaceus, pemphigoid, linear IgA disease, autoimmune haemolytic anaemia, Coombs positive haemolytic anaemia, acquired pernicious anaemia, juvenile pernicious anaemia, myalgic encephalitis/Royal Free Disease, chronic mucocutaneous candidiasis, giant cell arteritis,
- 20 primary sclerosing hepatitis, cryptogenic autoimmune hepatitis, Acquired Immunodeficiency Disease Syndrome, Acquired Immunodeficiency Related Diseases, Hepatitis C, common varied immunodeficiency (common variable hypogammaglobulinaemia), dilated cardiomyopathy, female infertility, ovarian failure, premature ovarian failure, fibrotic lung disease, cryptogenic fibrosing alveolitis, post-
- 25 inflammatory interstitial lung disease, interstitial pneumonitis, connective tissue disease associated interstitial lung disease, mixed connective tissue disease associated lung disease, systemic sclerosis associated interstitial lung disease, rheumatoid arthritis associated interstitial lung disease, systemic lupus erythematosus associated lung disease, dermatomyositis/polymyositis associated lung disease, Sjögren's disease
- 30 associated lung disease, ankylosing spondylitis associated lung disease, vasculitic diffuse lung disease, haemosiderosis associated lung disease, drug-induced interstitial

- lung disease, radiation fibrosis, bronchiolitis obliterans, chronic eosinophilic pneumonia, lymphocytic infiltrative lung disease, postinfectious interstitial lung disease, gouty arthritis, autoimmune hepatitis, type-1 autoimmune hepatitis (classical autoimmune or lupoid hepatitis), type-2 autoimmune hepatitis (anti-LKM antibody hepatitis),
- 5 autoimmune mediated hypoglycaemia, type B insulin resistance with acanthosis nigricans, hypoparathyroidism, acute immune disease associated with organ transplantation, chronic immune disease associated with organ transplantation, osteoarthritis, primary sclerosing cholangitis, psoriasis type 1, psoriasis type 2, idiopathic leucopaenia, autoimmune neutropaenia, renal disease NOS,
- 10 glomerulonephritides, microscopic vasculitis of the kidneys, Lyme disease, discoid lupus erythematosus, male infertility idiopathic or NOS, sperm autoimmunity, multiple sclerosis (all subtypes), sympathetic ophthalmia, pulmonary hypertension secondary to connective tissue disease, Goodpasture's syndrome, pulmonary manifestation of polyarteritis nodosa, acute rheumatic fever, rheumatoid spondylitis, Still's disease,
- 15 systemic sclerosis, Sjögren's syndrome, Takayasu's disease/arteritis, autoimmune thrombocytopaenia, idiopathic thrombocytopaenia, autoimmune thyroid disease, hyperthyroidism, goitrous autoimmune hypothyroidism (Hashimoto's disease), atrophic autoimmune hypothyroidism, primary myxoedema, phacogenic uveitis, primary vasculitis and vitiligo. The human antibodies, and antibody portions of the invention can
- 20 be used to treat humans suffering from autoimmune diseases, in particular those associated with inflammation, including, rheumatoid spondylitis, allergy, autoimmune diabetes, autoimmune uveitis, acute liver disease, chronic liver diseases, allergy and asthma, mental disorders (*e.g.*, depression and schizophrenia), and Th2 Type and Th1 Type mediated diseases.
- 25 Preferably, the antibodies of the invention or antigen-binding portions thereof, are used to treat rheumatoid arthritis, Crohn's disease, multiple sclerosis, insulin dependent diabetes, mellitus, and psoriasis.

An antibody, or antibody portion, of the invention also can be administered with one or more additional therapeutic agents useful in the treatment of autoimmune and

30 inflammatory diseases.

- 25 -

Antibodies of the invention, or antigen binding portions thereof can be used alone or in combination to treat such diseases. It should be understood that the antibodies of the invention or antigen binding portion thereof can be used alone or in combination with an additional agent, *e.g.*, a therapeutic agent, said additional agent
5 being selected by the skilled artisan for its intended purpose. For example, the additional agent can be a therapeutic agent art-recognized as being useful to treat the disease or condition being treated by the antibody of the present invention. The additional agent also can be an agent which imparts a beneficial attribute to the therapeutic composition *e.g.*, an agent which effects the viscosity of the composition.

10 It should further be understood that the combinations which are to be included within this invention are those combinations useful for their intended purpose. The agents set forth below are illustrative for purposes and not intended to be limited. The combinations which are part of this invention can be the antibodies of the present invention and at least one additional agent selected from the lists below. The
15 combination can also include more than one additional agent, *e.g.*, two or three additional agents if the combination is such that the formed composition can perform its intended function.

Preferred combinations are non-steroidal anti-inflammatory drug(s) also referred to as NSAIDS which include drugs like ibuprofen. Other preferred combinations are
20 corticosteroids including prednisolone; the well known side-effects of steroid use can be reduced or even eliminated by tapering the steroid dose required when treating patients in combination with the anti-IL-18 antibodies of this invention. Non-limiting examples of therapeutic agents for rheumatoid arthritis with which an antibody, or antibody
portion, of the invention can be combined include the following: cytokine suppressive
25 anti-inflammatory drug(s) (CSAIDs); antibodies to or antagonists of other human cytokines or growth factors, for example, TNF, LT, IL-1, IL-2, IL-6, IL-7, IL-8, IL-12, IL-15, IL-16, EMAP-II, GM-CSF, FGF, and PDGF. Antibodies of the invention, or antigen binding portions thereof, can be combined with antibodies to cell surface
molecules such as CD2, CD3, CD4, CD8, CD25, CD28, CD30, CD40, CD45, CD69,
30 CD80 (B7.1), CD86 (B7.2), CD90, or their ligands including CD154 (gp39 or CD40L).

Preferred combinations of therapeutic agents may interfere at different points in the autoimmune and subsequent inflammatory cascade; preferred examples include TNF antagonists like chimeric, humanized or human TNF antibodies, D2E7, (PCT Publication No. WO 97/29131), CA2 (RemicadeTM), CDP 571, and soluble p55 or p75 TNF receptors, derivatives, thereof, (p75TNFR1gG (EnbrelTM) or p55TNFR1gG (Lenercept), and also TNF α converting enzyme (TACE) inhibitors; similarly IL-1 inhibitors (Interleukin-1-converting enzyme inhibitors, IL-1RA *etc.*) may be effective for the same reason. Other preferred combinations include Interleukin 11. Yet another preferred combination are other key players of the autoimmune response which may act parallel to, dependent on or in concert with IL-18 function; especially preferred are IL-12 antagonists including IL-12 antibodies or soluble IL-12 receptors, or IL-12 binding proteins. It has been shown that IL-12 and IL-18 have overlapping but distinct functions and a combination of antagonists to both may be most effective. Yet another preferred combination are non-depleting anti-CD4 inhibitors. Yet other preferred combinations include antagonists of the co-stimulatory pathway CD80 (B7.1) or CD86 (B7.2) including antibodies, soluble receptors or antagonistic ligands.

The antibodies of the invention, or antigen binding portions thereof, may also be combined with agents, such as methotrexate, 6-MP, azathioprine sulphasalazine, mesalazine, olsalazine chloroquine/hydroxychloroquine, pencillamine, aurothiomalate (intramuscular and oral), azathioprine, cochlincine, corticosteroids (oral, inhaled and local injection), beta-2 adrenoreceptor agonists (salbutamol, terbutaline, salmeteral), xanthines (theophylline, aminophylline), cromoglycate, nedocromil, ketotifen, ipratropium and oxitropium, cyclosporin, FK506, rapamycin, mycophenolate mofetil, leflunomide, NSAIDs, for example, ibuprofen, cox-2 inhibitors, cox-2 selective inhibitors (*e.g.*, rofecoxib (VIOXXTM; Merck & Co., Inc.)) corticosteroids such as prednisolone, phosphodiesterase inhibitors, adenosine agonists, antithrombotic agents, complement inhibitors, adrenergic agents, agents which interfere with signalling by proinflammatory cytokines such as TNF α or IL-1 (*e.g.* IRAK, NIK, IKK, p38 or MAP kinase inhibitors), IL-1 β converting enzyme inhibitors, TNF α converting enzyme (TACE) inhibitors, T-cell signalling inhibitors such as kinase inhibitors, metalloproteinase inhibitors,

- 27 -

sulfasalazine, azathioprine, 6-mercaptopurines, angiotensin converting enzyme inhibitors, soluble cytokine receptors and derivatives thereof (e.g. soluble p55 or p75 TNF receptors and the derivatives p75TNFRIgG (Enbrel™ and p55TNFRIgG (Lenercept)), sIL-1RI, sIL-1RII, sIL-6R) and antiinflammatory cytokines (e.g. IL-4, IL-10, IL-11, IL-13 and TGFβ). Preferred combinations include methotrexate or leflunomide and in moderate or severe rheumatoid arthritis cases, cyclosporine.

Non-limiting examples of therapeutic agents for inflammatory bowel disease with which an antibody, or antibody portion, of the invention can be combined include the following: budenoside; epidermal growth factor; corticosteroids; cyclosporin, sulfasalazine; aminosalicylates; 6-mercaptopurine; azathioprine; metronidazole; lipoxygenase inhibitors; mesalamine; olsalazine; balsalazide; antioxidants; thromboxane inhibitors; IL-1 receptor antagonists; anti-IL-1β monoclonal antibodies; anti-IL-6 monoclonal antibodies; growth factors; elastase inhibitors; pyridinyl-imidazole compounds; antibodies to or antagonists of other human cytokines or growth factors, for example, TNF, LT, IL-1, IL-2, IL-6, IL-7, IL-8, IL-12, IL-15, IL-16, EMAP-II, GM-CSF, FGF, and PDGF. Antibodies of the invention, or antigen binding portions thereof, can be combined with antibodies to cell surface molecules such as CD2, CD3, CD4, CD8, CD25, CD28, CD30, CD40, CD45, CD69, CD90 or their ligands. The antibodies of the invention, or antigen binding portions thereof, may also be combined with agents, such as methotrexate, cyclosporin, FK506, rapamycin, mycophenolate mofetil, leflunomide, NSAIDs, for example, ibuprofen, corticosteroids such as prednisolone, phosphodiesterase inhibitors, adenosine agonists, antithrombotic agents, complement inhibitors, adrenergic agents, agents which interfere with signalling by proinflammatory cytokines such as TNFα or IL-1 (e.g. IRAK, NIK, IKK, p38 or MAP kinase inhibitors), IL-1β converting enzyme inhibitors, TNFα converting enzyme inhibitors, T-cell signalling inhibitors such as kinase inhibitors, metalloproteinase inhibitors, sulfasalazine, azathioprine, 6-mercaptopurines, angiotensin converting enzyme inhibitors, soluble cytokine receptors and derivatives thereof (e.g. soluble p55 or p75 TNF receptors, sIL-1RI, sIL-1RII, sIL-6R) and antiinflammatory cytokines (e.g. IL-4, IL-10, IL-11, IL-13 and TGFβ).

Preferred examples of therapeutic agents for Crohn's disease in which an antibody or an antigen binding portion can be combined include the following: TNF antagonists, for example, anti-TNF antibodies, D2E7 (PCT Publication No. WO 97/29131), CA2 (RemicadeTM), CDP 571, TNFR-Ig constructs, (p75TNFRIgG (EnbrelTM) and p55TNFRIgG (Lenercept)) inhibitors and PDE4 inhibitors. Antibodies of the invention or antigen binding portions thereof, can be combined with corticosteroids, for example, budenoside and dexamethasone. Antibodies of the invention or antigen binding portions thereof, may also be combined with agents such as sulfasalazine, 5-aminosalicylic acid and olsalazine, and agents which interfere with synthesis or action of proinflammatory cytokines such as IL-1, for example, IL-1 β converting enzyme inhibitors and IL-1ra. Antibodies of the invention or antigen binding portion thereof may also be used with T cell signaling inhibitors, for example, tyrosine kinase inhibitors 6-mercaptopurines. Antibodies of the invention or antigen binding portions thereof, can be combined with IL-11.

Non-limiting examples of therapeutic agents for multiple sclerosis with which an antibody, or antibody portion, of the invention can be combined include the following: corticosteroids; prednisolone; methylprednisolone; azathioprine; cyclophosphamide; cyclosporine; methotrexate; 4-aminopyridine; tizanidine; interferon- β 1a (Avonex; Biogen); interferon- β 1b (Betaseron; Chiron/Berlex); Copolymer 1 (Cop-1; Copaxone; Teva Pharmaceutical Industries, Inc.); hyperbaric oxygen; intravenous immunoglobulin; clabribine; antibodies to or antagonists of other human cytokines or growth factors, for example, TNF, LT, IL-1, IL-2, IL-6, IL-7, IL-8, IL-12, IL-15, IL-16, EMAP-II, GM-CSF, FGF, and PDGF. Antibodies of the invention, or antigen binding portions thereof, can be combined with antibodies to cell surface molecules such as CD2, CD3, CD4, CD8, CD25, CD28, CD30, CD40, CD45, CD69, CD80, CD86, CD90 or their ligands. The antibodies of the invention, or antigen binding portions thereof, may also be combined with agents, such as methotrexate, cyclosporine, FK506, rapamycin, mycophenolate mofetil, leflunomide, NSAIDs, for example, ibuprofen, corticosteroids such as prednisolone, phosphodiesterase inhibitors, adenosine agonists, antithrombotic agents, complement inhibitors, adrenergic agents, agents which interfere with signalling

- 29 -

by proinflammatory cytokines such as TNF α or IL-1 (*e.g.* IRAK, NIK, IKK, p38 or MAP kinase inhibitors), IL-1 β converting enzyme inhibitors, TACE inhibitors, T-cell signalling inhibitors such as kinase inhibitors, metalloproteinase inhibitors, sulfasalazine, azathioprine, 6-mercaptopurines, angiotensin converting enzyme inhibitors, soluble cytokine receptors and derivatives thereof (*e.g.* soluble p55 or p75 TNF receptors, sIL-1RI, sIL-1RII, sIL-6R) and antiinflammatory cytokines (*e.g.* IL-4, IL-10, IL-13 and TGF β).

Preferred examples of therapeutic agents for multiple sclerosis in which the antibody or antigen binding portion thereof can be combined to include interferon- β , for example, IFN β 1a and IFN β 1b; copaxone, corticosteroids, IL-1 inhibitors, TNF inhibitors, and antibodies to CD40 ligand and CD80.

The pharmaceutical compositions of the invention may include a "therapeutically effective amount" or a "prophylactically effective amount" of an antibody or antibody portion of the invention. A "therapeutically effective amount" refers to an amount effective, at dosages and for periods of time necessary, to achieve the desired therapeutic result. A therapeutically effective amount of the antibody or antibody portion may vary according to factors such as the disease state, age, sex, and weight of the individual, and the ability of the antibody or antibody portion to elicit a desired response in the individual. A therapeutically effective amount is also one in which any toxic or detrimental effects of the antibody or antibody portion are outweighed by the therapeutically beneficial effects. A "prophylactically effective amount" refers to an amount effective, at dosages and for periods of time necessary, to achieve the desired prophylactic result. Typically, since a prophylactic dose is used in subjects prior to or at an earlier stage of disease, the prophylactically effective amount will be less than the therapeutically effective amount.

Dosage regimens may be adjusted to provide the optimum desired response (*e.g.*, a therapeutic or prophylactic response). For example, a single bolus may be administered, several divided doses may be administered over time or the dose may be proportionally reduced or increased as indicated by the exigencies of the therapeutic situation. It is especially advantageous to formulate parenteral compositions in dosage

unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein refers to physically discrete units suited as unitary dosages for the mammalian subjects to be treated; each unit containing a predetermined quantity of active compound calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. The specification for the dosage unit forms of the invention are dictated by and directly dependent on (a) the unique characteristics of the active compound and the particular therapeutic or prophylactic effect to be achieved, and (b) the limitations inherent in the art of compounding such an active compound for the treatment of sensitivity in individuals.

10 An exemplary, non-limiting range for a therapeutically or prophylactically effective amount of an antibody or antibody portion of the invention is 0.1-20 mg/kg, more preferably 1-10 mg/kg. It is to be noted that dosage values may vary with the type and severity of the condition to be alleviated. It is to be further understood that for any particular subject, specific dosage regimens should be adjusted over time according to the individual need and the professional judgment of the person administering or supervising the administration of the compositions, and that dosage ranges set forth herein are exemplary only and are not intended to limit the scope or practice of the claimed composition.

20 Methods of Making Anti-IL-18 Antibodies

The anti-IL-18 antibodies of the invention are made using any one of a variety of techniques known in the art for preparing antibodies and using antigens comprising the IL-18 peptide epitope described in subsection I, *i.e.*, an epitope of human IL-18 comprising amino acids PLFEDMTDSDCRDNA (SEQ ID NO: 1).

25 In general, the methods of the invention for making an antibody that binds human interleukin-18 (IL-18) involve:

 exposing an antibody repertoire to an antigen comprising an epitope of human IL-18 comprising amino acids PLFEDMTDSDCRDNA (SEQ ID NO: 1), or portion thereof (*e.g.*, SEQ ID NO: 3 or 33); and

selecting from the antibody repertoire an antibody that binds the epitope of human IL-18 comprising amino acids PLFEDMTDSDCRDNA (SEQ ID NO: 1), or portion thereof (*e.g.*, SEQ ID NO: 3 or 33).

In one embodiment, the antibody repertoire is an *in vivo* repertoire in an animal
5 and the method comprises immunizing the animal with the antigen comprising the epitope of human IL-18 comprising amino acids PLFEDMTDSDCRDNA (SEQ ID NO: 1). In another embodiment, the antibody repertoire is a recombinant antibody library and the method comprising screening the library with the antigen comprising the epitope of human IL-18 comprising amino acids PLFEDMTDSDCRDNA (SEQ ID NO: 1).
10 Preferably, the library is a human antibody library.

Methods for immunizing an animal with an antigen to thereby raise specific antibodies to the antigen are well known in the art. An IL-18 antigen comprising an epitope of human IL-18 comprising amino acids PLFEDMTDSDCRDNA (SEQ ID NO: 1) can be administered to an animal to elicit polyclonal antibodies and specific
15 antibodies that bind the epitope can be isolated by selecting from the polyclonal antibodies those antibodies that bind to the epitope (*e.g.*, by passing the polyclonal antisera over a column that comprises a peptide comprising amino acids PLFEDMTDSDCRDNA (SEQ ID NO: 1) of hIL-18). The antigen used to elicit the polyclonal antibodies can be intact (*i.e.*, full-length) hIL-18 or can be a portion of hIL-18
20 that includes the epitope of interest, *e.g.*, a synthetic peptide comprising amino acids PLFEDMTDSDCRDNA (SEQ ID NO: 1) of hIL-18. Furthermore, monoclonal antibodies to the epitope can be made from the aforementioned animals using standard hybridoma technology and selection for those hybridomas secreting an antibody that specifically binds the epitope of interest, *e.g.*, by screening the hybridomas with a
25 peptide comprising amino acids PLFEDMTDSDCRDNA (SEQ ID NO: 1) of hIL-18 and selecting for antibodies that bind specifically to the peptide.

In vitro methods also can be used to make the antibodies of the invention, wherein an antibody library is screened to identify an antibody having the desired binding specificity. Methods for such screening of recombinant antibody libraries are
30 well known in the art and include methods described in, for example, Ladner *et al.* U.S. Patent No. 5,223,409; Kang *et al.* PCT Publication No. WO 92/18619; Dower *et al.* PCT

Publication No. WO 91/17271; Winter *et al.* PCT Publication No. WO 92/20791; Markland *et al.* PCT Publication No. WO 92/15679; Breitling *et al.* PCT Publication No. WO 93/01288; McCafferty *et al.* PCT Publication No. WO 92/01047; Garrard *et al.* PCT Publication No. WO 92/09690; Fuchs *et al.* (1991) *Bio/Technology* 9:1370-1372; 5 Hay *et al.* (1992) *Hum Antibod Hybridomas* 3:81-85; Huse *et al.* (1989) *Science* 246:1275-1281; McCafferty *et al.*, *Nature* (1990) 348:552-554; Griffiths *et al.* (1993) *EMBO J* 12:725-734; Hawkins *et al.* (1992) *J Mol Biol* 226:889-896; Clackson *et al.* (1991) *Nature* 352:624-628; Gram *et al.* (1992) *PNAS* 89:3576-3580; Garrad *et al.* (1991) *Bio/Technology* 9:1373-1377; Hoogenboom *et al.* (1991) *Nuc Acid Res* 19:4133- 10 4137; and Barbas *et al.* (1991) *PNAS* 88:7978-7982, and PCT Publication No. WO 97/29131, the contents of each of which are incorporated herein by reference.

The recombinant antibody library may be from a subject immunized with IL-18, or a portion of IL-18 comprising the epitope of amino acids PLFEDMTDSDCRDNA (SEQ ID NO: 1). Alternatively, the recombinant antibody library may be from a naïve 15 subject, *i.e.*, one who has not been immunized with IL-18, such as a human antibody library from a human subject who has not been immunized with human IL-18. Antibodies of the invention are selected by screening the recombinant antibody library with the epitope of amino acids PLFEDMTDSDCRDNA (SEQ ID NO: 1) of human IL-18 to thereby select those antibodies that recognize this epitope. Methods for 20 conducting such screening and selection are well known in the art, such as described in the references in the preceding paragraph.

To select antibodies of the invention having a particular binding affinity for hIL-18, the art-known method of surface plasmon resonance can be used. To select antibodies having a particular neutralizing activity for hIL-18, standard methods known 25 in the art for assessing the inhibition of hIL-18 activity may be used. In addition, methods for immunizing mice that have been transgenically altered to encode a human immunoglobulin repertoire thereby enabling the organism to express fully human antibodies in response to an immunogen, are known in the art (see, *e.g.*, U.S.P.N.s 5,877,397 and 6,150,584).

Uses of Anti-IL-18 Antibodies

Given their ability to bind to hIL-18, the anti-hIL-18 antibodies, or portions thereof, of the invention can be used to detect hIL-18 (*e.g.*, in a biological sample, such as serum or plasma), using a conventional immunoassay, such as an enzyme linked
5 immunosorbent assays (ELISA), an radioimmunoassay (RIA) or tissue immunohistochemistry. The invention provides a method for detecting hIL-18 in a biological sample comprising contacting a biological sample with an antibody, or antibody portion, of the invention and detecting either the antibody (or antibody portion) bound to hIL-18 or unbound antibody (or antibody portion), to thereby detect hIL-18 in
10 the biological sample. The antibody is directly or indirectly labeled with a detectable substance to facilitate detection of the bound or unbound antibody. Suitable detectable substances include various enzymes, prosthetic groups, fluorescent materials, luminescent materials and radioactive materials. Examples of suitable enzymes include horseradish peroxidase, alkaline phosphatase, β -galactosidase, or acetylcholinesterase;
15 examples of suitable prosthetic group complexes include streptavidin/biotin and avidin/biotin; examples of suitable fluorescent materials include umbelliferone, fluorescein, fluorescein isothiocyanate, rhodamine, dichlorotriazinylamine fluorescein, dansyl chloride or phycoerythrin; an example of a luminescent material includes luminol; and examples of suitable radioactive material include ^{125}I , ^{131}I , ^{35}S , ^{32}P ,
20 ^{33}P , or ^3H .

Alternative to labeling the antibody, hIL-18 can be assayed in biological fluids by a competition immunoassay utilizing rhIL-18 standards labeled with a detectable substance and an unlabeled anti-hIL-18 antibody. In this assay, the biological sample, the labeled rhIL-18 standards and the anti-hIL-18 antibody are combined and the amount
25 of labeled rhIL-18 standard bound to the unlabeled antibody is determined. The amount of hIL-18 in the biological sample is inversely proportional to the amount of labeled rhIL-18 standard bound to the anti-hIL-18 antibody.

The antibodies and antibody portions of the invention preferably are capable of neutralizing hIL-18 activity both *in vitro* and *in vivo*. Accordingly, such antibodies and
30 antibody portions of the invention can be used to inhibit hIL-18 activity, *e.g.*, in a cell culture containing hIL-18, in human subjects or in other mammalian subjects having IL-

18 with which an antibody of the invention cross-reacts. In one embodiment, the invention provides a method for inhibiting IL-18 activity comprising contacting IL-18 with an antibody or antibody portion of the invention such that IL-18 activity is inhibited. Preferably, the IL-18 is human IL-18. For example, in a cell culture
5 containing, or suspected of containing hIL-18, an antibody or antibody portion of the invention can be added to the culture medium to inhibit hIL-18 activity in the culture.

In another embodiment, the invention provides a method for inhibiting IL-18 activity in a subject suffering from a disorder in which IL-18 activity is detrimental. The invention provides methods for inhibiting IL-18 activity in a subject suffering from such
10 a disorder, which method comprises administering to the subject an antibody or antibody portion of the invention such that IL-18 activity in the subject is inhibited. Preferably, the IL-18 is human IL-18 and the subject is a human subject. Alternatively, the subject can be a mammal expressing an IL-18 with which an antibody of the invention cross-reacts. Still further the subject can be a mammal into which has been introduced hIL-18
15 (*e.g.*, by administration of hIL-18 or by expression of an hIL-18 transgene). An antibody of the invention can be administered to a human subject for therapeutic purposes. Moreover, an antibody of the invention can be administered to a non-human mammal expressing an IL-18 with which the antibody cross-reacts for veterinary purposes or as an animal model of human disease. Regarding the latter, such animal
20 models may be useful for evaluating the therapeutic efficacy of antibodies of the invention (*e.g.*, testing of dosages and time courses of administration).

In particular, one animal model for modulating IL-18 activity in an animal uses NOD-SCID mice which are transplanted with human peripheral blood mononuclear cells. Then, two to four weeks after engraftment (as measured by human IgG titers in
25 serum) the mice are injected with LPS (lipopolysaccharide). Four to six hours later LPS-induced human interferon-gamma serum titers are determined. The efficacy (potency) of anti-IL-18 antibodies (*e.g.*, IL-18 neutralizing antibodies) is determined by injecting the antibodies (ip) one day prior to LPS challenge followed by monitoring the test animals for a reduction in interferon-gamma serum titers (a function of IL-18 *in vivo*
30 activity) as compared to controls (see, *e.g.*, Holmes et al., Hybridoma, 19:363367 (2000)).

- 35 -

As used herein, the term "a disorder in which IL-18 activity is detrimental" is intended to include diseases and other disorders in which the presence of IL-18 in a subject suffering from the disorder has been shown to be, or is suspected of being, either responsible for the pathophysiology of the disorder or a factor that contributes to a worsening of the disorder. Accordingly, a disorder in which IL-18 activity is detrimental is a disorder in which inhibition of IL-18 activity is expected to alleviate the symptoms and/or progression of the disorder. Such disorders may be evidenced, for example, by an increase in the concentration of IL-18 in a biological fluid of a subject suffering from the disorder (*e.g.*, an increase in the concentration of IL-18 in serum, plasma, synovial fluid, *etc.* of the subject), which can be detected, for example, using an anti-IL-18 antibody as described above.

Non-limiting examples of disorders that can be treated with the antibodies of the invention include those disorders discussed in the section above pertaining to pharmaceutical compositions of the antibodies of the invention.

Other features of the invention will be apparent from the following examples which should not be construed as limiting.

EXEMPLIFICATION

In general, the practice of the present invention employs, unless otherwise indicated, conventional techniques of chemistry, molecular biology, recombinant DNA technology, PCR technology, immunology (especially, *e.g.*, antibody technology), and any necessary cell culture or animal husbandry techniques, which are within the skill of the art and are explained fully in the literature. See, *e.g.*, Sambrook, Fritsch and Maniatis, *Molecular Cloning: Cold Spring Harbor Laboratory Press* (1989); *DNA Cloning*, Vols. 1 and 2, (D.N. Glover, Ed. 1985); *Oligonucleotide Synthesis* (M.J. Gait, Ed. 1984); *PCR Handbook Current Protocols in Nucleic Acid Chemistry*, Beaucage, Ed. John Wiley & Sons (1999) (Editor); *Oxford Handbook of Nucleic Acid Structure*, Neidle, Ed., Oxford Univ Press (1999); *PCR Protocols: A Guide to Methods and Applications*, Innis *et al.*, Academic Press (1990); *PCR Essential Techniques: Essential Techniques*, Burke, Ed., John Wiley & Son Ltd (1996); *The PCR Technique: RT-PCR*, Siebert, Ed., Eaton Pub. Co. (1998); *Quantitative PCR Protocols*, Kochanowski *et al.*,

Eds., Humana Press (1999); *Clinical Applications of PCR*, Lo, Ed., Humana Press (1998); *Antibody Engineering Protocols (Methods in Molecular Biology)*, 510, Paul, S., Humana Pr (1996); *Antibody Engineering: A Practical Approach (Practical Approach Series, 169)*, McCafferty, Ed., Irl Pr (1996); *Antibodies: A Laboratory Manual*, Harlow
5 *et al.*, C.S.H.L. Press, Pub. (1999); *Current Protocols in Molecular Biology*, eds. Ausubel *et al.*, John Wiley & Sons (1992); *Large-Scale Mammalian Cell Culture Technology*, Lubiniecki, A., Ed., Marcel Dekker, Pub., (1990); and *Manipulating the Mouse Embryo*, Hogan *et al.*, C.S.H.L. Press, Pub (1994).

Throughout the examples, unless otherwise indicated, the above materials and
10 methods were used.

EXAMPLE 1

ISOLATION OF ANTI-IL-18 ANTIBODIES

Antibodies to hIL-18 were isolated by screening separate scFv phage display
15 libraries prepared using human VL and VH cDNAs from mRNA derived from human B cells (*e.g.*, tonsils and spleen). Construction of the library and methods for selection are described in Vaughan *et al.* (1996) *Nature Biotech.* 14: 309-314.

The libraries were screened using either full length human IL-18 (SEQ ID NO: 61), a peptide epitope of IL-18 (SEQ ID NOS: 1-3), or a panel of overlapping 15 amino
20 acid peptides representing IL-18 (the epitope sequence of which is presented in Table 5; SEQ ID NOS: 31-60). IL-18 specific antibodies were selected by coating the antigen onto immunotubes using standard procedures (Marks *et al.*, (1991) *J. Mol. Biol.* 222: 581-597). The scFv libraries were screened using either IL-18, a peptide epitope of IL-18, or an IL-18 peptide panel to generate a significant number of IL-18 specific binders.
25 Several different clonotypes were selected, determined by restriction enzyme digestion patterns, and confirmed by DNA sequencing.

In order to identify IL-18 antibodies which preferentially bind either full length IL-18 or a representative peptide thereof, the supernatant containing scFv was titrated on biotin-captured IL-18 in an ELISA and binding characteristics were determined.

Two anti-IL-18 single chain antibodies were obtained, one termed 2E1, independently isolated using a peptide epitope and the peptide panel, and a second anti-IL-18 antibody termed LT28, isolated using full length IL-18. These parent anti-IL-18 antibodies were selected for further study and modification.

5

EXAMPLE 2

AFFINITY MATURATION OF AN ANTI-18 ANTIBODIES

A single chain Fv version of antibody 2E1 having an identified IL-18 binding activity and the heavy chain and light chain sequence shown in Table 6 was further
10 modified for improved neutralization of IL-18 activity.

Table 6. Sequence of Single-Chain Anti-IL-18 Antibody 2E1

2E1 Heavy Chain (SEQ ID NO: 18)	
CDR1 (SEQ ID NO: 9) QVQLVQSGAEVKKPGASMKVSCKTSGYTF <u>FTGYI</u> HWVRQAHGQGFEWI	
CDR2 (SEQ ID NO: 10) <u>GRLNPTTGDANFAEK</u> FQGRVALTRDTSISTAYLQLDSLKSDDTAVYYCAG	CDR3 (SEQ ID NO: 11) <u>KEGA</u> WGQG
TLVTVSS	

Table 6. Sequence of Single-Chain Anti-IL-18 Antibody 2E1 (Continued)

2E1 Light Chain (SEQ ID NO: 19)	
CDR1 (SEQ ID NO: 12)	CDR2 (SEQ ID NO: 13)
SSELTQDPAVSVALGQTVRITC <u>QGDSL</u> RHFYPNWWYQQKPGQAPVLVIY <u>GKNNRPS</u>	
CDR3 (SEQ ID NO: 14)	
GIPDRFSGSGSGNTGSLTITGAQAEDEADYYC <u>GSRDSSGIHVV</u> FGGGTKVTVLG	

The anti-IL-18 antibody 2E1 was independently selected using an IL-18 peptide
5 and sequential, overlapping, peptide panel representative of IL-18 (see Table 6).

The specific amino acid residues of the heavy chain variable region selected for
mutagenesis are summarized in Table 7. In particular, with respect to the heavy chain
region, individual amino acid substitutions were tested at positions H30, H31, H32, H33,
and H35 of CDR1, positions H52, H52a, H53, H54, H56, and H58 of CDR2, and H95,
10 H96, H97, and H98 of CDR3.

With regards to light chain amino acid residues selected for mutagenesis,
individual amino acid substitutions were tested at positions L30, L31, L32, and L34 of
CDR1, positions L50, L52, L53, and L55 of CDR2 and positions L89, L90, L91, L92,
L93, L94, L95, L95a, L95b, L96, and L97 of CDR 3.

Table 7. Heavy Chain Amino Acid Substitutions Introduced Into 2E1

Heavy Chain Mutations	
CDR / Kabat Position	substituted residue
CDR1	
H30	A, R, N, D, C, G, H, I, F, P, S, or V
H31	A, C, H, S, T, or Y
H32	R, N, C, H, P, S, or T
H33	N, D, C, Q, H, L, M, F, S, or V
H35	N, D, L, or F
CDR2	
H52	T
H52a	R, Q, L, S, T, or W
H53	A, R, N, L, P, S, or Y
H54	A, R, N, D, Q, L, K, M, P, S, or Y
H56	A, R, N, C, G, H, I, L, or F
H58	A, R, Q, E, H, I, L, K, M, F, S, T, Y, P, S, T, W, Y, or V
CDR3	
H95	A, R, E, Q, S, Y, V, H, P, W, or C
H96	A, R, Q, S, Y, V, H, P, W, or C
H97	A, R, E, Q, S, Y, V, H, P, W, or C
H98	R, E, Q, S, Y, V, H, P, W, or C

Table 8. Light Chain Amino Acid Substitutions Introduced Into 2E1

5

Light Chain Mutations	
CDR / Kabat Position	substituted residue
CDR1	

- 40 -

L30	N, D, C, G, I, L, S, W, or Y
L31	R, N, D, C, G, H, I, L, P, S, T, or Y
L32	R, N, D, E, G, I, L, P, S, T, or V
L34	A, R, N, D, E, H, I, L, K, M, F, P, S, T, Y, or V
CDR2	
L50	A, N, I, L, F, P, S, W, Y, or V
L52	A, R, D, E, H, I, L, M, F, P, S, T, or V
L53	A, R, C, I, L, K, M, P, S, or T
L55	A, R, N, D, C, G, H, I, L, S, T, or Y
CDR3	
L89	A, R, E, Q, S, Y, V, H, P, W, or C
L90	A, R, E, Q, Y, V, H, P, W, or C
L91	R, E, Q, S, Y, V, H, P, W, or C
L92	A, R, E, Q, S, Y, V, H, P, W, or C
L93	A, R, E, Q, Y, V, H, P, W, or C
L94	A, R, E, Q, Y, V, H, P, W, or C
L95	A, R, E, Q, S, Y, V, H, P, W, or C
L95a	A, R, E, Q, S, Y, V, H, P, W, or C
L95b	A, R, E, Q, S, Y, V, P, W, or C
L96	A, R, E, Q, S, Y, H, P, W, or C
L97	A, R, E, Q, S, Y, H, P, W, or C

Substitutions were introduced using standard techniques (*e.g.*, as described in Taylor et al., *Nucleic Acids Res.* 13: 8764-8758 (1985); Nakamaye et al., *Nucleic Acids Res.* 14: 9679-9698 (1986); and Olsen et al., *Methods in Enzymology*, 217: 189 (1993)).

- 5 In brief, oligonucleotides degenerate for a given codon were synthesized for each of the positions to be mutagenized. A single-stranded DNA template was prepared from the original plasmid containing a single-chain Fv version of the antibody 2E1 gene. The nucleic acid sequence of the parent 2E1 antibody heavy and light chain is provided in SEQ ID NOS: 62 and 64. The mutant oligonucleotides were then used to create a

complementary DNA strand and eventually a double-stranded plasmid, thus incorporating the degeneracy or the different mutations in a given codon of the antibody. In particular, the CDR3 region of the heavy and light chain of 2E1 was altered using the QuikChange Kit (Stratagene) according to the manufactures instruction.

- 5 A representative number of clones were then sequenced from each mutagenesis reaction (*i.e.*, 7 to 36 clones) and those representing a change from the parent 2E1 single chain antibody sequence were expressed in bacteria and purified for further *in vitro* and *in vivo* testing as described infra.

- 10 In another screen using a full length IL-18 ligand, a second anti-IL-18 antibody was identified and selected for further improvement using affinity maturation. In particular, using the techniques described above, the LT28 antibody having the heavy chain and light chain sequence shown in Table 9 (and nucleic sequence provided in SEQ ID NOS: 66 and 68) was further modified.

15 **Table 9. Sequence of Single-Chain Anti-IL-18 Antibody LT28**

LT28 Heavy Chain (SEQ ID NO: 28)	
CDR1 (SEQ ID NO: 20)	CDR2 (SEQ ID NO: 21)
LVQPGGSLRLSCAASGFTFS	<u>SYAMS</u> WVRQAPGKGLEWVS <u>AI</u> <u>SGSGGSTYYADSVKG</u>
CDR3 (SEQ ID NO: 22)	
RFTISRDN SKNTLYLQMNSLRAEDTAVYYCAR	<u>DDDDYDFDY</u> WGRGTMVTVSS

Table 9. Sequence of Single-Chain Anti-IL-18 Antibody LT28 (Continued)

LT28 Light Chain (SEQ ID NO: 29)	
CDR1 (SEQ ID NO: 23)	CDR2 (SEQ ID NO: 24)
QSVLTQPPSASGTPGQRVTISC <u>SGSSSNIGINAVN</u> WYQQLPGTAPKLLIY <u>GNDQRPS</u>	
CDR3 (SEQ ID NO: 25)	
GVPDRFSGSKSGTSASLAISGLQSEDEADYYC <u>AAWDDSLSGPV</u> FGGGTKLTVLG	

With respect to the heavy chain region, amino acid substitutions were introduced at positions H31, H32, H33, and H35 of CDR1, positions H50, H51, H52, H52a, H53, H54, H56, and H58 of CDR2, and H95, H96, H97, H98, H99, H100, H100a, H101, and H102 of CDR3.

- 5 With regards to light chain residues selected for mutation, amino acid substitutions were introduced at positions L30, L31, L32, L34 of CDR1, positions L50, L52, L53, L55 of CDR2 and positions L89, L90, L91, L92, L93, L94, L95, L95a, L95b, L96, L97.

10 **Table 10. Heavy Chain Amino Acid Substitutions Introduced into LT28**

Heavy Chain Mutations	
CDR / Kabat Position	substituted residue
CDR1	
H31	A, R, E, Q, S, Y, V, H, P, W, or C
H32	A, R, E, Q, S, Y, V, H, P, W, or C
H33	A, R, E, Q, S, Y, V, H, P, W, or C
H35	A, R, E, Q, S, Y, V, H, P, W, or C
CDR2	
H50	A, R, E, Q, S, Y, V, H, P, W, or C
H51	A, R, E, Q, S, Y, V, H, P, W, or C
H52	A, R, E, Q, S, Y, V, H, P, W, or C
H52a	A, R, E, Q, S, Y, V, H, P, W, or C
H53	A, R, E, Q, S, Y, V, H, P, W, or C
H54	A, R, E, Q, S, Y, V, H, P, W, or C
H56	A, R, E, Q, S, Y, V, H, P, W, or C
H58	A, R, E, Q, S, Y, V, H, P, W, C
CDR3	
H95	A, R, E, Q, S, Y, V, H, P, W, or C

- 44 -

H96	A, R, E, Q, S, Y, V, H, P, W, or C
H97	A, R, E, Q, S, Y, V, H, P, W, or C
H98	A, R, E, Q, S, Y, V, H, P, W, or C
H99	A, R, E, Q, S, Y, V, H, P, W, or C
H100	A, R, E, Q, S, Y, V, H, P, W, or C
H100a	A, R, E, Q, S, Y, V, H, P, W, or C
H101	A, R, E, Q, S, Y, V, H, P, W, or C
H102	A, R, E, Q, S, Y, V, H, P, W, or C

Table 11. Light Chain Amino Acid Substitutions Introduced into LT28

Light Chain Mutations	
CDR / Position	substituted residue
CDR1	
L30	A, R, E, Q, S, Y, V, H, P, W, C
L31	A, R, E, Q, S, Y, V, H, P, W, C
L32	R, E, Q, S, Y, V, H, P, W, C, G
L34	A, R, E, Q, S, Y, V, H, P, W, C
CDR2	
L50	A, R, E, Q, S, Y, V, H, P, C
L52	A, R, E, S, Y, V, H, P, W, C
L53	A, R, E, S, Y, V, H, P, W, C, N
L55	A, E, Q, S, Y, V, H, P, W, C
CDR3	
L89	A, R, E, Q, S, Y, V, H, P, W, C
L90	A, R, E, Q, S, Y, V, H, P, W, C
L91	A, R, E, Q, S, Y, V, H, P, W, C
L92	A, R, E, Q, S, Y, V, H, P, W, C
L93	A, R, E, Q, S, Y, V, H, P, W, C
L94	A, R, E, Q, S, Y, V, H, P, W, C
L95	A, R, E, Q, S, Y, V, H, P, W, C
L95a	A, R, E, Q, S, Y, V, H, P, W, C
L95b	A, R, E, Q, S, Y, V, H, P, W, C
L96	A, R, E, Q, S, Y, V, H, P, W, C
L97	A, R, E, Q, S, Y, V, H, P, W, C

Substitutions were introduced as described above. A representative number of
5 clones were then sequenced from each mutagenesis reaction and those representing a

change from the parent LT28 single chain antibody sequence were expressed in bacteria and purified for further testing as described below.

EXAMPLE 3

5 BINDING ACTIVITY OF HUMAN ANTIBODIES TO IL-18

Real-time binding interactions between ligand (biotinylated recombinant human IL-18 (rhIL-18) immobilized on a biosensor matrix) and analyte (antibodies in solution) were measured by surface plasmon resonance (SPR) using the BIAcore system (Pharmacia Biosensor, Piscataway, NJ). The system utilizes the optical properties of
10 SPR to detect alterations in protein concentrations within a dextran biosensor matrix. Proteins are covalently bound to the dextran matrix at known concentrations. Antibodies are injected through the dextran matrix and specific binding between injected antibodies and immobilized ligand results in an increased matrix protein concentration and resultant change in the SPR signal. These changes in SPR signal are recorded as
15 resonance units (RU) and are displayed with respect to time along the y-axis of a sensorgram.

To facilitate immobilization of biotinylated rhIL-18 on the biosensor matrix, streptavidin is covalently linked via free amine groups to the dextran matrix by first activating carboxyl groups on the matrix with 100 mM N-hydroxysuccinimide (NHS)
20 and 400 mM N-ethyl-N'-(3-diethylaminopropyl) carbodiimide hydrochloride (EDC). Next, streptavidin is injected across the activated matrix. Thirty-five microliters of streptavidin (25 µg/ml), diluted in sodium acetate, pH 4.5, is injected across the activated biosensor and free amines on the protein are bound directly to the activated carboxyl groups. Unreacted matrix EDC-esters are deactivated by an injection of 1 M
25 ethanolamine. Streptavidin-coupled biosensor chips also are commercially available (Pharmacia BR-1000-16, Pharmacia Biosensor, Piscataway, NJ).

Biotinylated rhIL-18 was prepared by first dissolving 5.0 mg of biotin (D-biotinyl-ε-aminocaproic acid N-hydroxysuccinimide ester; Boehringer Mannheim Cat. No. 1008 960) in 500 µl dimethylsulfoxide to make a 10 mg/ml solution. Ten
30 microliters of biotin was added per ml of rhIL-18 (at 2.65 mg/ml) for a 2:1 molar ratio of

- 47 -

biotin to rhIL-18. The reaction was mixed gently and incubated for two hours at room temperature in the dark. A PD-10 column, Sephadex G-25M (Pharmacia Catalog No. 17-0851-01) was equilibrated with 25 ml of cold PBS and loaded with 2 ml of rhIL-18-biotin per column. The column was eluted with 10 x 1 ml cold PBS. Fractions were
5 collected and read at OD280 (1.0 OD = 1.25 mg/ml). The appropriate fractions were pooled and stored at -80° C until use.

Biotinylated rhIL-18 to be immobilized on the matrix via streptavidin was diluted in PBS running buffer (Gibco Cat. No. 14190-144, Gibco BRL, Grand Island, NY) supplemented with 0.05% (BIAcore) surfactant P20 (Pharmacia BR-1000-54,
10 Pharmacia Biosensor, Piscataway, NJ). To determine the capacity of rhIL-18-specific antibodies to bind immobilized rhIL-18, a binding assay was conducted as follows. Aliquots of biotinylated rhIL-18 (25 nM; 10 μ l aliquots) were injected through the streptavidin-coupled dextran matrix at a flow rate of 5 μ l/min. Before injection of the protein and immediately afterward, PBS buffer alone flowed through each flow cell.
15 The net difference in signal between baseline and approximately 30 sec. after completion of biotinylated rhIL-18 injection was taken to represent the binding value. Direct rhIL-18-specific antibody binding to immobilized biotinylated rhIL-18 was measured. Antibodies (20 μ g/ml) were diluted in PBS running buffer and 25 μ l aliquots were injected through the immobilized protein matrices at a flow rate of 5 μ l/min. Prior
20 to injection of antibody, and immediately afterwards, PBS buffer alone flowed through each flow cell. The net difference in baseline signal and signal after completion of antibody injection was taken to represent the binding value of the particular sample. Biosensor matrices were regenerated using 100 mM HCl before injection of the next sample. To determine the off rate (K_{off}), on rate (K_{on}), association rate (K_a) and
25 dissociation rate (K_d) constants, BIAcore kinetic evaluation software (version 2.1) was used.

Representative results of improved candidate anti-IL-18 antibodies binding to biotinylated rhIL-18, as compared to the parent antibodies 2E1 and LT28 (and murine controls), are shown below in Table 12. For comparison, IC50 values from the cell-
30 based neutralization assay are also included and these are described in Example 4. All

clones were prepared as single-chain Fv antibodies for testing using Biacore analysis and the cell-based assay described below. Parental clones listed comprise an unmutated parental heavy and light chain, whereas single chain mutants contain one parental chain and one mutated chain where the mutated chain is indicated as being either heavy (H) or light (L) followed by the Kabat position and nature of the amino acid substitution.

Table 12. Binding of Anti-IL-18 Antibodies Derived From 2E1 and LT28

Antibody Clone	On-rates (M ⁻¹ s ⁻¹)	Off-rates (s ⁻¹)	Kd (M)	IC50 Value*
2E1 parent and mutants				
2E1 (parent) ScFv	2.6E+3	6.42E-03	1.5E-07	3.3E-8M
2E1 (parent) IgG				9.0E-10M
L34S		1.69E-04		1.5E-8M
H53R		2.34E-03		2.5E-8M
H53Y		-		1.5E-8M
H58Q		-		1.6E-8M
L34S + H53R (2E1RS)	2.7E+03	6.82E-05	2.3E-08	3.0E-09M
L34S + H58Q		-		1.5E-8M
L34S + H53Y		5.28E-05		6.7E-9M
H53R + H58Q		-		1.2E-8M
H53Y + H58Q		-		1.2E-8M
L34S + H53R + H58Q		6.18E-05		2.8E-9M
L34S + H53Y + H58Q		-		8.0E-9M
L90C				4x
L93C				2-4x

- 49 -

L94P, Q, or R				2-4x
L95R, Y				3-8x
L95bE, W				2-4x
LT28 parent and mutants				
LT28 (parent)	1.3E+04	4.8E-04	3.9E-08	9.0E-8M
H54Q				2-3x
H58W				2-3x
1				
125-2H	1.7E+05	1.1E-04	6.2E-10	2.E-10M
318-M	1.2E+04	1.1E-04	9.6E-09	4.0E-9M

* Some values presented as fold improvement compared to the parent.

EXAMPLE 4

NEUTALIZING ACTIVITY OF ANTI-IL-18 ANTIBODIES

To examine the neutralizing activity of the anti-human IL-18 antibodies of the invention, an art recognized assay for monitoring IL-18 activity was used.

5 Briefly, the assay employs KG1 cells (ATCC #CCL-246, myelogenous leukemia bone marrow cells) which were cultured according to standard techniques (*e.g.*, using RPMI 1640 Culture Medium Gibco #21870-076; (supplemented with 10% Fetal Bovine Serum (BioWhittaker #14-501F); 2mM L-glutamine (Gibco #25030-081); 50units/ml Penicillin, 50ug/ml Streptomycin (Gibco #15070-063); and .075% Sodium Bicarbonate).

10 To test for IL-18 neutralization, 3 x 10⁵ KG-1 cells stimulated with 20ng/ml hTNF-alpha (Lot# 19130132) was incubated with 50ul of anti-IL-18 antibody (4 x Conc.) and 50ul of IL-18 (4xConc.=8ng/ml) and incubated for 37°C for 1hr or 16-20 hrs. To determine the amount of IL-18 neutralization that occurred as a function of induced hIFN-gamma production, an ELISA was performed using commercially available Elisa
15 Kits (R & D #DIF00/Endogen #EH-IFNG), according to the manufacturers instructions, and hIFN-gamma production was calculated (pg/ml) off a standard curve.

In all, four mutant antibodies, *i.e.*, the 2E1 derived L34S, H53R, H53Y, and H58Q, were shown to have greater IL-18 neutralization potency than the parent 2E1 antibody (see Table 12). The improvements in IC50 values using the KG-1 assay were
20 in the range of 2 to 5 fold, and similar improved binding results were determined using BIAcore analysis.

Various mutation combination clones were also prepared and tested, and this data is summarized in the Table 12. The best combination clone L34S-H53R showed a greater than ten fold improvement over the parent antibody 2E1 in both the KG-1 cell-
25 based assay and using the BIAcore analysis. The resulting antibody was designated the name of 2E1RS.

Several other mutant clones of 2E1 showed an improvement in potency, *i.e.*, IL-18 neutralization, as determined using the KG-1 assay. The mutant L95Y offered 5 to 8 fold better IC50 values than the parent 2E1 antibody. Several other mutants offered a 2
30 to 3 fold improvement and they are 2E1 mutants H96A, H96Q, H96S, H98S, L90C,

- 51 -

L90W, L93C, L94P, L94P, L94Q, L94R, L94W, L95R, L95aA, L95aH, L95aP, L95aR, L95aW, L95bE, L95bW, L95bY, L97C, and L97E.

The binding of 2E1 in the form of an ScFv antibody or IgG antibody was also compared (see Fig. 5).

- 5 Still further, two mutants derived from the LT28 parent should improved 1L-18 neutralization activity compared to the parent antibody.

These results demonstrate that fully human IL-18 neutralizing antibodies can be obtained using the methods and compositions of the invention.

10

EQUIVALENTS

Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention described herein. Such equivalents are intended to be encompassed by the following claims.

What is claimed:

1. A compound capable of binding a human IL-18 amino acid sequence, or portion thereof, wherein said amino acid comprises a sequence selected from the group consisting of SEQ ID NO: 67 and SEQ ID NO: 68.
5
2. The compound of claim 1, wherein said compound is selected from the group consisting of a small molecule, peptide, polypeptide, antibody, and antibody fragment.
10
3. The compound of claim 2, wherein said antibody, or antibody fragment, is fully human.
4. A human monoclonal antibody, or antigen-binding portion thereof,
15 capable of binding to human IL-18.
5. The antibody of claim 4, wherein the antibody, or antigen-binding portion thereof, dissociates from human IL-18 with a k_{off} rate constant of 0.1s^{-1} or less, as determined by surface plasmon resonance, or which inhibits human IL-18 activity with
20 an IC_{50} of $1 \times 10^{-6}\text{M}$ or less.
6. The antibody of claim 4, wherein the antibody, or antigen-binding portion thereof, dissociates from human IL-18 with a k_{off} rate constant of $1 \times 10^{-2}\text{s}^{-1}$ or less, as determined by surface plasmon resonance, or which inhibits human IL-18 activity with
25 an IC_{50} of $1 \times 10^{-7}\text{M}$ or less.
7. The antibody of claim 4, wherein the antibody, or antigen-binding portion thereof, dissociates from human IL-18 with a k_{off} rate constant of $1 \times 10^{-3}\text{s}^{-1}$ or less, as determined by surface plasmon resonance, or which inhibits human IL-18 activity with
30 an IC_{50} of $1 \times 10^{-8}\text{M}$ or less.

8. The antibody of claim 4, wherein the antibody, or antigen-binding portion thereof, dissociates from human IL-18 with a k_{off} rate constant of $1 \times 10^{-4} \text{s}^{-1}$ or less, as determined by surface plasmon resonance, or which inhibits human IL-18 activity with an IC_{50} of $1 \times 10^{-9} \text{M}$ or less.

5

9. The antibody of claim 4, wherein the antibody, or antigen-binding portion thereof, dissociates from human IL-18 with a k_{off} rate constant of $1 \times 10^{-5} \text{s}^{-1}$ or less, as determined by surface plasmon resonance, or which inhibits human IL-18 activity with an IC_{50} of $1 \times 10^{-10} \text{M}$ or less.

10

10. The antibody of claim 4, wherein the antibody, or antigen-binding portion thereof, dissociates from human IL-18 with a k_{off} rate constant of $1 \times 10^{-6} \text{s}^{-1}$ or less, as determined by surface plasmon resonance, or which inhibits human IL-18 activity with an IC_{50} of $1 \times 10^{-11} \text{M}$ or less.

15

11. An isolated antibody, or an antigen-binding portion thereof, that binds an epitope of human IL-18, or portion thereof, comprising an amino acid sequence selected from the group comprising SEQ ID NO: 3 and SEQ ID NO: 33.

20

12. The antibody, or antigen-binding portion thereof, of claim 11, wherein the antibody is a neutralizing antibody.

13. The antibody, or antigen-binding portion thereof, of claim 11, which is a human antibody.

25

14. The antibody, or antigen-binding portion thereof, of claim 11, which is a recombinant antibody.

15. The antibody, or antigen-binding portion thereof, of claim 11, which is a
30 monoclonal antibody.

16. An isolated antibody, or antigen-binding portion thereof, that binds to an epitope of human IL-18, wherein the antibody, or antigen-binding portion thereof, dissociates from human IL-18 with a k_{off} rate constant of 0.1s^{-1} or less, as determined by surface plasmon resonance, or which inhibits human IL-18 activity with an IC_{50} of $1 \times 10^{-6}\text{M}$ or less.

17. The isolated antibody of claim 16, or an antigen-binding portion thereof, which dissociates from human IL-18 with a k_{off} rate constant of $1 \times 10^{-2}\text{s}^{-1}$ or less, as determined by surface plasmon resonance, or which inhibits human IL-18 activity with an IC_{50} of $1 \times 10^{-7}\text{M}$ or less.

18. The isolated antibody of claim 16, or an antigen-binding portion thereof, which dissociates from human IL-18 with a k_{off} rate constant of $1 \times 10^{-3}\text{s}^{-1}$ or less, as determined by surface plasmon resonance, or which inhibits human IL-18 activity with an IC_{50} of $1 \times 10^{-8}\text{M}$ or less.

19. The isolated antibody of claim 16, or an antigen-binding portion thereof, which dissociates from human IL-18 with a k_{off} rate constant of $1 \times 10^{-4}\text{s}^{-1}$ or less, as determined by surface plasmon resonance, or which inhibits human IL-18 activity with an IC_{50} of $1 \times 10^{-9}\text{M}$ or less.

20. The isolated antibody of claim 16, or an antigen-binding portion thereof, which dissociates from human IL-18 with a k_{off} rate constant of $1 \times 10^{-5}\text{s}^{-1}$ or less, as determined by surface plasmon resonance, or which inhibits human IL-18 activity with an IC_{50} of $1 \times 10^{-10}\text{M}$ or less.

21. The isolated antibody of claim 16, or an antigen-binding portion thereof, which dissociates from human IL-18 with a k_{off} rate constant of $1 \times 10^{-6}\text{s}^{-1}$ or less, as determined by surface plasmon resonance, or which inhibits human IL-18 activity with an IC_{50} of $1 \times 10^{-11}\text{M}$ or less.

- 55 -

22. An isolated human antibody, or an antigen-binding portion thereof, comprising at least one variable region CDR domain capable of binding an epitope of human IL-18.

5 23. The isolated antibody, or an antigen-binding portion thereof, of claim 22, wherein said antibody, or an antigen-binding portion thereof, contains at least one amino acid substitution or insertion that improves IL-18 binding as compared to the unmodified antibody or antigen-binding portion thereof.

10 24. The isolated antibody, or an antigen-binding portion thereof, of claim 22, wherein said antibody, or an antigen-binding portion thereof, contains at least one amino acid substitution or insertion that improves neutralization of IL-18 as compared to the unmodified antibody or antigen-binding portion thereof.

15 25. The isolated antibody, or an antigen-binding portion thereof, of claim 22, wherein said variable region comprises a CDR domain selected from the group consisting of:

 a heavy chain CDR1 domain having an amino acid sequence of SEQ ID NO: 9, or sequence modified from SEQ ID NO: 9 by at least one amino acid substitution;

20 a heavy chain CDR2 domain having an amino acid sequence of SEQ ID NO: 10, or sequence modified from SEQ ID NO: 10 by at least one amino acid substitution; and

 a heavy chain CDR3 domain having an amino acid sequence of SEQ ID NO: 11, or sequence modified from SEQ ID NO: 11 by at least one amino acid substitution.

25 26. The isolated antibody, or an antigen-binding portion thereof, of claim 25, wherein said variable region comprises a CDR domain selected from the group consisting of:

 a heavy chain CDR1 domain modified from SEQ ID NO: 9 by at least one amino acid substitution at position H30, H31, H32, H33, or H35;

30 a heavy chain CDR2 domain modified from SEQ ID NO: 10 by at least one amino acid substitution at position H52, H52a, H53, H54, H56, or H58; and

- 56 -

a heavy chain CDR3 domain modified from SEQ ID NO: 11 by at least one amino acid substitution at position H95, H96, H97, or H98.

27. The isolated antibody, or an antigen-binding portion thereof, of claim 22,
5 wherein said variable region comprises a CDR domain selected from the group consisting of:

a light chain CDR1 domain having an amino acid sequence of SEQ ID NO: 12 ,
or sequence modified from SEQ ID NO: 12 by at least one amino acid substitution;

a light chain CDR2 domain having an amino acid sequence of SEQ ID NO: 13 ,
10 or sequence modified from SEQ ID NO: 13 by at least one amino acid substitution; and

a light chain CDR3 domain having an amino acid sequence of SEQ ID NO: 14 ,
or sequence modified from SEQ ID NO: 14 by at least one amino acid substitution.

28. The isolated antibody, or an antigen-binding portion thereof, of claim 27,
15 wherein said variable region comprises a CDR domain selected from the group consisting of:

a light chain CDR1 domain modified from SEQ ID NO: 12 by at least one amino
acid substitution at position L30, L31, L32, or L34;

a light chain CDR2 domain modified from SEQ ID NO: 13 by at least one amino
20 acid substitution at position L50, L52, L53, or L55; and

a light chain CDR3 domain modified from SEQ ID NO: 14 by at least one amino
acid substitution at position L89, L90, L91, L92, L93, L94, L95, L95a, L95b, L96, or
L97.

25 29. An isolated antibody, or an antigen-binding portion thereof, with a variable
region comprising an amino acid sequence selected from the group consisting of SEQ ID
NO: 15, 16, and 17.

- 57 -

30. An isolated antibody, or an antigen-binding portion thereof, with a light chain variable region (LCVR) comprising the amino acid sequence of SEQ ID NO: 15 and a heavy chain variable region (HCVR) comprising the amino acid sequence of SEQ ID NO: 16.

5

31. An isolated antibody, or an antigen-binding portion thereof, with a light chain variable region (LCVR) having the amino acid sequence of SEQ ID NO: 15 and a heavy chain variable region (HCVR) having the amino acid sequence of SEQ ID NO: 17.

10

32. The isolated antibody, or an antigen-binding portion thereof, of claim 22, wherein said variable region comprises a CDR domain selected from the group consisting of:

- a heavy chain CDR1 domain having an amino acid sequence of SEQ ID NO: 20 ,
- 15 or sequence modified from SEQ ID NO: 20 by at least one amino acid substitution;
- a heavy chain CDR2 domain having an amino acid sequence of SEQ ID NO: 21 ,
- or sequence modified from SEQ ID NO: 21 by at least one amino acid substitution; and
- a heavy chain CDR3 domain having an amino acid sequence of SEQ ID NO: 22 ,
- or sequence modified from SEQ ID NO: 22 by at least one amino acid substitution.

20

33. The isolated antibody, or an antigen-binding portion thereof, of claim 32, wherein said variable region comprises a CDR domain selected from the group consisting of:

- a heavy chain CDR1 domain modified from SEQ ID NO: 20 by at least one
- 25 amino acid substitution at position H30, H31, H32, H33, or H35;
- a heavy chain CDR2 domain modified from SEQ ID NO: 21 by at least one amino acid substitution at position H50, H51, H52, H52a, H53, H54, H56, or H58; and
- a heavy chain CDR3 domain modified from SEQ ID NO: 22 by at least one amino acid substitution at position H96, H96, H97, H98, H99, H100, H100a, H101, or
- 30 H102.

34. The isolated antibody, or an antigen-binding portion thereof, of claim 32, wherein said variable region comprises a CDR domain selected from the group consisting of:

5 a light chain CDR1 domain having an amino acid sequence of SEQ ID NO: 23 , or sequence modified from SEQ ID NO: 23 by at least one amino acid substitution at position ;

a light chain CDR2 domain having an amino acid sequence of SEQ ID NO: 24 , or sequence modified from SEQ ID NO: 24 by at least one amino acid substitution; and

10 a light chain CDR3 domain having an amino acid sequence of SEQ ID NO: 25, or sequence modified from SEQ ID NO: 25 by at least one amino acid substitution.

35. The isolated antibody, or an antigen-binding portion thereof, of claim 34, wherein said variable region comprises a CDR domain selected from the group consisting of:

15 a light chain CDR1 domain modified from SEQ ID NO: 23 by at least one amino acid substitution at position L30, L31, L32, or L34;

a light chain CDR2 domain modified from SEQ ID NO: 24 by at least one amino acid substitution at position L50, L52, L53, or L55; and

20 a light chain CDR3 domain modified from SEQ ID NO: 25 by at least one amino acid substitution at position L89, L90, L91, L92, L93, L94, L95, L95a, L95b, L96, or L97.

36. An isolated antibody, or an antigen-binding portion thereof, with a variable region comprising an amino acid selected from the group consisting of SEQ ID NO: 26, 27, and 29.

25

37. An isolated antibody, or an antigen-binding portion thereof, with a light chain variable region (LCVR) comprising the amino acid sequence of SEQ ID NO: 29 and a heavy chain variable region (HCVR) comprising the amino acid sequence of SEQ ID NO: 26.

30

- 59 -

38. An isolated antibody, or an antigen-binding portion thereof, with a light chain variable region (LCVR) having the amino acid sequence of SEQ ID NO: 29 and a heavy chain variable region (HCVR) having the amino acid sequence of SEQ ID NO: 27.

5

39. An isolated nucleic acid encoding an antibody CDR amino acid sequence of any one of claims 4-38.

40. The isolated nucleic acid of claim 39, which is in a recombinant expression vector.

10

41. A host cell into which the recombinant expression vector of claim 40 has been introduced.

15

42. A method of synthesizing an antibody that binds human IL-18, comprising culturing the host cell of claim 41 in a culture medium until an antibody that binds human IL-18 is synthesized by the cell.

20

43. The method of claim 42, wherein said antibody is human.

44. A pharmaceutical composition comprising the antibody, or antigen-binding portion thereof, of any of claims 4-38, and a pharmaceutically acceptable carrier.

25

45. The pharmaceutical composition of claim 44 which further comprises at least one additional therapeutic agent for treating a disorder in which IL-18 activity is detrimental.

- 60 -

46. The pharmaceutical composition of claim 45, wherein said additional agent is selected from the group consisting of an antibody, or fragment thereof, capable of binding human IL-12, methotrexate anti-TNF, corticosteroids, cyclosporin, rapamycin, FK506, and non-steroidal anti-inflammatory agents.

5

47. A method of making an antibody that binds human interleukin-18 (IL-18), comprising:

exposing an antibody repertoire to an antigen comprising an epitope of human IL-18 or portion thereof; and

10 selecting from the antibody repertoire an antibody that binds the epitope of human IL-18, or portion thereof.

48. The method of claim 47, wherein the antibody repertoire is an *in vivo* repertoire in an animal and the method comprises immunizing the animal with the
15 antigen comprising an epitope of human IL-18 or portion thereof.

49. The method of claim 46 or 47, wherein said epitope comprises the amino acid sequence selected from the group consisting of SEQ ID NO: 3 and 33.

20 50. The method of claim 47, wherein said *in vivo* repertoire is a fully human immunoglobulin repertoire integrated into the genome of the animal.

51. The method of claim 47, wherein the antibody repertoire is a recombinant antibody library and the method comprises screening the library with an antigen
25 comprising the epitope of human IL-18 or portion thereof.

52. The method of claim 47, wherein the library is a human antibody library.

53. A method for inhibiting human IL-18 activity comprising contacting
30 human IL-18 with the compound of claim 1 such that human IL-18 activity is inhibited.

- 61 -

54. A method for inhibiting human IL-18 activity comprising contacting human IL-18 with the antibody, or antigen-binding portion thereof, of any of claims 4-38 such that human IL-18 activity is inhibited.

5 55. A method for inhibiting human IL-18 activity in a human subject suffering from a disorder in which IL-18 activity is detrimental, comprising administering to the human subject the compound of claim 1 such that human IL-18 activity in the human subject is inhibited.

10 56. A method for inhibiting human IL-18 activity in a human subject suffering from a disorder in which IL-18 activity is detrimental, comprising administering to the human subject the antibody, or antigen-binding portion thereof, of any of claims 4-38 such that human IL-18 activity in the human subject is inhibited.

15 57. A method for treating a human subject suffering from a disorder in which IL-18 activity is detrimental by administering a compound according to claim 1 such that treatment is achieved.

58. A method for treating a human subject suffering from a disorder in which
20 IL-18 activity is detrimental by administering an antibody according to any one of claims 4-38 such that treatment is achieved.

59. The method of claim 57 or 58, wherein said disorder is selected from the group comprising rheumatoid arthritis, osteoarthritis, juvenile chronic arthritis, Lyme
25 arthritis, psoriatic arthritis, reactive arthritis, spondyloarthropathy, systemic lupus erythematosus, Crohn's disease, ulcerative colitis, inflammatory bowel disease, insulin dependent diabetes mellitus, thyroiditis, asthma, allergic diseases, psoriasis, dermatitis scleroderma, graft versus host disease, organ transplant rejection, acute or chronic immune disease associated with organ transplantation, sarcoidosis, atherosclerosis,
30 disseminated intravascular coagulation, Kawasaki's disease, Grave's disease, nephrotic syndrome, chronic fatigue syndrome, Wegener's granulomatosis, Henoch-Schoenlein


- purpura, microscopic vasculitis of the kidneys, chronic active hepatitis, uveitis, septic shock, toxic shock syndrome, sepsis syndrome, cachexia, infectious diseases, parasitic diseases, acquired immunodeficiency syndrome, acute transverse myelitis, Huntington's chorea, Parkinson's disease, Alzheimer's disease, stroke, primary biliary cirrhosis,
- 5 hemolytic anemia, malignancies, heart failure, myocardial infarction, Addison's disease, sporadic, polyglandular deficiency type I and polyglandular deficiency type II, Schmidt's syndrome, adult (acute) respiratory distress syndrome, alopecia, alopecia areata, seronegative arthropathy, arthropathy, Reiter's disease, psoriatic arthropathy, ulcerative colitic arthropathy, enteropathic synovitis, chlamydia, yersinia and salmonella
- 10 associated arthropathy, spondyloarthropathy, atheromatous disease/arteriosclerosis, atopic allergy, autoimmune bullous disease, pemphigus vulgaris, pemphigus foliaceus, pemphigoid, linear IgA disease, autoimmune haemolytic anaemia, Coombs positive haemolytic anaemia, acquired pernicious anaemia, juvenile pernicious anaemia, myalgic encephalitis/Royal Free Disease, chronic mucocutaneous candidiasis, giant cell arteritis,
- 15 primary sclerosing hepatitis, cryptogenic autoimmune hepatitis, Acquired Immunodeficiency Disease Syndrome, Acquired Immunodeficiency Related Diseases, Hepatitis C, common varied immunodeficiency (common variable hypogammaglobulinaemia), dilated cardiomyopathy, female infertility, ovarian failure, premature ovarian failure, fibrotic lung disease, cryptogenic fibrosing alveolitis, post-
- 20 inflammatory interstitial lung disease, interstitial pneumonitis, connective tissue disease associated interstitial lung disease, mixed connective tissue disease associated lung disease, systemic sclerosis associated interstitial lung disease, rheumatoid arthritis associated interstitial lung disease, systemic lupus erythematosus associated lung disease, dermatomyositis/polymyositis associated lung disease, Sjögren's disease
- 25 associated lung disease, ankylosing spondylitis associated lung disease, vasculitic diffuse lung disease, haemosiderosis associated lung disease, drug-induced interstitial lung disease, radiation fibrosis, bronchiolitis obliterans, chronic eosinophilic pneumonia, lymphocytic infiltrative lung disease, postinfectious interstitial lung disease, gouty arthritis, autoimmune hepatitis, type-1 autoimmune hepatitis (classical autoimmune or
- 30 lupoid hepatitis), type-2 autoimmune hepatitis (anti-LKM antibody hepatitis), autoimmune mediated hypoglycaemia, type B insulin resistance with acanthosis

- nigricans, hypoparathyroidism, acute immune disease associated with organ transplantation, chronic immune disease associated with organ transplantation, osteoarthritis, primary sclerosing cholangitis, psoriasis type 1, psoriasis type 2, idiopathic leucopaenia, autoimmune neutropaenia, renal disease NOS,
- 5 glomerulonephritides, microscopic vasculitis of the kidneys, Lyme disease, discoid lupus erythematosus, male infertility idiopathic or NOS, sperm autoimmunity, multiple sclerosis (all subtypes), sympathetic ophthalmia, pulmonary hypertension secondary to connective tissue disease, Goodpasture's syndrome, pulmonary manifestation of polyarteritis nodosa, acute rheumatic fever, rheumatoid spondylitis, Still's disease,
- 10 systemic sclerosis, Sjögren's syndrome, Takayasu's disease/arteritis, autoimmune thrombocytopaenia, idiopathic thrombocytopaenia, autoimmune thyroid disease, hyperthyroidism, goitrous autoimmune hypothyroidism (Hashimoto's disease), atrophic autoimmune hypothyroidism, primary myxoedema, phacogenic uveitis, primary vasculitis, vitiligo, acute liver disease, chronic liver diseases, allergy and asthma, mental
- 15 disorders (*e.g.*, depression and schizophrenia) and Th2 Type and Th1 Type mediated diseases.

60. A method of treating a patient suffering from a disorder in which IL-18 is detrimental comprising the step of administering an anti-IL-18 antibody, before,
- 20 concurrent, or after the administration of a second agent, wherein the second agent is selected from the group consisting of an anti-IL-12 antibody or antigen binding fragment thereof, methotrexate, anti-TNF antibody or antigen binding fragment thereof, corticosteroids, cyclosporin, rapamycin, FK506, and non-steroidal anti-inflammatory agents.


1/5

Fig. 1




2/5

Fig. 2




3/5

Fig. 3




4/5

Fig. 4



5/5

Fig. 5



SEQUENCE LISTING

<110> Ghayur, Tarig et al.

<120> ANTIBODIES THAT BIND HUMAN INTERLEUKIN-18 AND METHODS
OF MAKING AND USING

<130> BBI-149

<140>

<141>

<150> 60/181,608

<151> 2000-02-10

<160> 71

<170> PatentIn Ver. 2.1

<210> 1

<211> 15

<212> PRT

<213> Homo sapiens

<400> 1

Pro Leu Phe Glu Asp Met Thr Asp Ser Asp Cys Arg Asp Asn Ala
1 5 10 15

<210> 2

<211> 16

<212> PRT

<213> Homo sapiens

<400> 2

Cys Pro Leu Phe Glu Asp Met Thr Asp Ser Asp Cys Arg Asp Asn Ala
1 5 10 15

<210> 3

<211> 12

<212> PRT

<213> Homo sapiens

<400> 3

Pro Leu Phe Glu Asp Met Thr Asp Ser Asp Cys Arg
1 5 10

<210> 4

<211> 157

<212> PRT

<213> Homo sapiens

<400> 4

Tyr Phe Gly Lys Leu Glu Ser Lys Leu Ser Val Ile Arg Asn Leu Asn
1 5 10 15

Asp Gln Val Leu Phe Ile Asp Gln Gly Asn Arg Pro Leu Phe Glu Asp
20 25 30

Met Thr Asp Ser Asp Cys Arg Asp Asn Ala Pro Arg Thr Ile Phe Ile
 35 40 45

Ile Ser Met Tyr Lys Asp Ser Gln Pro Arg Gly Met Ala Val Thr Ile
 50 55 60

Ser Val Lys Cys Glu Lys Ile Ser Thr Leu Ser Cys Glu Asn Lys Ile
 65 70 75 80

Ile Ser Phe Lys Glu Met Asn Pro Pro Asp Asn Ile Lys Asp Thr Lys
 85 90 95

Ser Asp Ile Ile Phe Phe Gln Arg Ser Val Pro Gly His Asp Asn Lys
 100 105 110

Met Gln Phe Glu Ser Ser Ser Tyr Glu Gly Tyr Phe Leu Ala Cys Glu
 115 120 125

Lys Glu Arg Asp Leu Phe Lys Leu Ile Leu Lys Lys Glu Asp Glu Leu
 130 135 140

Gly Asp Arg Ser Ile Met Phe Thr Val Gln Asn Glu Asp
 145 150 155

<210> 5
 <211> 153
 <212> PRT
 <213> Homo sapiens

<400> 5
 Ala Pro Val Arg Ser Leu Asn Cys Thr Leu Arg Asp Ser Gln Gln Lys
 1 5 10 15

Ser Leu Val Met Ser Gly Pro Tyr Glu Leu Lys Ala Leu His Leu Gln
 20 25 30

Gly Gln Asp Met Glu Gln Gln Val Val Phe Ser Met Ser Phe Val Gln
 35 40 45

Gly Glu Glu Ser Asn Asp Lys Ile Pro Val Ala Leu Gly Leu Lys Glu
 50 55 60

Lys Asn Leu Tyr Leu Ser Cys Val Leu Lys Asp Asp Lys Pro Thr Leu
 65 70 75 80

Gln Leu Glu Ser Val Asp Pro Lys Asn Tyr Pro Lys Lys Lys Met Glu
 85 90 95

Lys Arg Phe Val Phe Asn Lys Ile Glu Ile Asn Asn Lys Leu Glu Phe
 100 105 110

Glu Ser Ala Gln Phe Pro Asn Trp Tyr Ile Ser Thr Ser Gln Ala Glu
 115 120 125

Asn Met Pro Val Phe Leu Gly Gly Thr Lys Gly Gly Gln Asp Ile Thr
 130 135 140

Asp Phe Thr Met Gln Phe Val Ser Ser
 145 150

<210> 6
 <211> 145
 <212> PRT
 <213> Homo sapiens

<400> 6
 Ser Ser Lys Met Gln Ala Phe Arg Ile Trp Asp Val Asn Gln Lys Thr
 1 5 10 15
 Phe Tyr Leu Arg Asn Asn Gln Leu Val Ala Gly Tyr Leu Gln Gly Pro
 20 25 30
 Asn Val Asn Leu Glu Glu Lys Ile Asp Val Val Pro Ile Glu Pro His
 35 40 45
 Ala Leu Phe Leu Gly Ile His Gly Gly Lys Met Cys Leu Ser Cys Val
 50 55 60
 Lys Ser Gly Asp Glu Thr Arg Leu Gln Leu Glu Ala Val Asn Ile Thr
 65 70 75 80
 Asp Leu Ser Glu Asn Arg Lys Gln Asp Lys Arg Phe Ala Phe Ile Arg
 85 90 95
 Ser Asp Ser Gly Pro Thr Thr Ser Phe Glu Ser Ala Ala Cys Pro Gly
 100 105 110
 Trp Phe Leu Cys Thr Ala Met Glu Ala Asp Gln Pro Val Ser Leu Thr
 115 120 125
 Asn Met Pro Asp Glu Gly Val Met Val Thr Lys Phe Tyr Phe Gln Glu
 130 135 140
 Asp
 145

<210> 7
 <211> 297
 <212> PRT
 <213> Homo sapiens

<400> 7
 Cys Thr Ser Arg Pro His Ile Thr Val Val Glu Gly Glu Pro Phe Tyr
 1 5 10 15
 Leu Lys His Cys Ser Cys Ser Leu Ala His Glu Ile Glu Thr Thr Thr
 20 25 30
 Lys Ser Trp Tyr Lys Ser Ser Gly Ser Gln Glu His Val Glu Leu Asn
 35 40 45
 Pro Arg Ser Ser Ser Arg Ile Ala Leu His Asp Cys Val Leu Glu Phe
 50 55 60
 Trp Pro Val Glu Leu Asn Asp Thr Gly Ser Tyr Phe Phe Gln Met Lys
 65 70 75 80
 Asn Tyr Thr Gln Lys Trp Lys Leu Asn Val Ile Arg Arg Asn Lys His

4
85 90 95

Ser Cys Phe Thr Glu Arg Gln Val Thr Ser Lys Ile Val Glu Val Lys
100 105 110

Lys Phe Phe Gln Ile Thr Cys Glu Asn Ser Tyr Tyr Gln Thr Leu Val
115 120 125

Asn Ser Thr Ser Leu Tyr Lys Asn Cys Lys Lys Leu Leu Leu Glu Asn
130 135 140

Asn Lys Asn Pro Thr Ile Lys Lys Asn Ala Glu Phe Glu Asp Gln Gly
145 150 155 160

Tyr Tyr Ser Cys Val His Phe Leu His His Asn Gly Lys Leu Phe Asn
165 170 175

Ile Thr Lys Thr Phe Asn Ile Thr Ile Val Glu Asp Arg Ser Asn Ile
180 185 190

Val Pro Val Leu Leu Gly Pro Lys Leu Asn His Val Ala Val Glu Leu
195 200 205

Gly Lys Asn Val Arg Leu Asn Cys Ser Ala Leu Leu Asn Glu Glu Asp
210 215 220

Val Ile Tyr Trp Met Phe Gly Glu Glu Asn Gly Ser Asp Pro Asn Ile
225 230 235 240

His Glu Glu Lys Glu Met Arg Ile Met Thr Pro Glu Gly Lys Trp His
245 250 255

Ala Ser Lys Val Leu Arg Ile Glu Asn Ile Gly Glu Ser Asn Leu Asn
260 265 270

Val Leu Tyr Asn Cys Thr Val Ala Ser Thr Gly Gly Thr Asp Thr Lys
275 280 285

Ser Phe Ile Leu Val Arg Lys Ala Asp
290 295

<210> 8
<211> 310
<212> PRT
<213> Homo sapiens

<400> 8
Cys Lys Glu Arg Glu Glu Lys Ile Ile Leu Val Ser Ser Ala Asn Glu
1 5 10 15

Ile Asp Val Arg Pro Cys Pro Leu Asn Pro Asn Glu His Lys Gly Thr
20 25 30

Ile Thr Trp Tyr Lys Asp Asp Ser Lys Thr Pro Val Ser Thr Glu Gln
35 40 45

Ala Ser Arg Ile His Gln His Lys Glu Lys Leu Trp Phe Val Pro Ala
50 55 60

Lys Val Glu Asp Ser Gly His Tyr Tyr Cys Val Val Arg Asn Ser Ser

```
<210> 9
<211> 6
<212> PRT
<213> Homo sapiens

<400> 9
Thr Gly Tyr Tyr Ile His
  1                               5
```

BNSDOCID: <WO 0158956A2.1 >

<213> Homo sapiens

<400> 10

Gly Arg Leu Asn Pro Thr Thr Gly Asp Ala Asn Phe Ala Glu Lys Phe
1 5 10 15

Gln

<210> 11

<211> 4

<212> PRT

<213> Homo sapiens

<400> 11

Lys Glu Gly Ala
1

<210> 12

<211> 11

<212> PRT

<213> Homo sapiens

<400> 12

Gln Gly Asp Ser Leu Arg His Phe Tyr Pro Asn
1 5 10

<210> 13

<211> 7

<212> PRT

<213> Homo sapiens

<400> 13

Gly Lys Asn Asn Arg Pro Ser
1 5

<210> 14

<211> 11

<212> PRT

<213> Homo sapiens

<400> 14

Gly Ser Arg Asp Ser Ser Gly Ile His Val Val
1 5 10

<210> 15

<211> 11

<212> PRT

<213> Homo sapiens

<400> 15

Gln Gly Asp Ser Leu Arg His Phe Tyr Ser Asn
1 5 10

<210> 16

<211> 17
 <212> PRT
 <213> Homo sapiens

<400> 16
 Gly Arg Leu Asn Pro Arg Thr Gly Asp Ala Asn Phe Ala Glu Lys Phe
 1 5 10 15

Gln

<210> 17
 <211> 17
 <212> PRT
 <213> Homo sapiens

<400> 17
 Gly Arg Leu Asn Pro Arg Thr Gly Asp Ala Gln Phe Ala Glu Lys Phe
 1 5 10 15

Gln

<210> 18
 <211> 113
 <212> PRT
 <213> Homo sapiens

<400> 18
 Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
 1 5 10 15

Ser Met Lys Val Ser Cys Lys Thr Ser Gly Tyr Thr Phe Thr Gly Tyr
 20 25 30

Tyr Ile His Trp Val Arg Gln Ala His Gly Gln Gly Phe Glu Trp Ile
 35 40 45

Gly Arg Leu Asn Pro Thr Thr Gly Asp Ala Asn Phe Ala Glu Lys Phe
 50 55 60

Gln Gly Arg Val Ala Leu Thr Arg Asp Thr Ser Ile Ser Thr Ala Tyr
 65 70 75 80

Leu Gln Leu Asp Ser Leu Lys Ser Asp Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

Ala Gly Lys Glu Gly Ala Trp Gly Gln Gly Thr Leu Val Thr Val Ser
 100 105 110

Ser

<210> 19
 <211> 109
 <212> PRT
 <213> Homo sapiens

<400> 19

Ser Ser Glu Leu Thr Gln Asp Pro Ala Val Ser Val Ala Leu Gly Gln
 1 5 10 15

Thr Val Arg Ile Thr Cys Gln Gly Asp Ser Leu Arg His Phe Tyr Pro
 20 25 30

Asn Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Val Leu Val Ile Tyr
 35 40 45

Gly Lys Asn Asn Arg Pro Ser Gly Ile Pro Asp Arg Phe Ser Gly Ser
 50 55 60

Gly Ser Gly Asn Thr Gly Ser Leu Thr Ile Thr Gly Ala Gln Ala Glu
 65 70 75 80

Asp Glu Ala Asp Tyr Tyr Cys Gly Ser Arg Asp Ser Ser Gly Ile His
 85 90 95

Val Val Phe Gly Gly Gly Thr Lys Val Thr Val Leu Gly
 100 105

<210> 20

<211> 4

<212> PRT

<213> Homo sapiens

<400> 20

Ser Tyr Ala Met
 1

<210> 21

<211> 17

<212> PRT

<213> Homo sapiens

<400> 21

Ala Ile Ser Gly Ser Gly Gly Ser Thr Tyr Tyr Ala Asp Ser Val Lys
 1 5 10 15

Gly

<210> 22

<211> 9

<212> PRT

<213> Homo sapiens

<400> 22

Asp Asp Asp Tyr Asp Phe Asp Tyr
 1 5

<210> 23

<211> 13

<212> PRT

<213> Homo sapiens

<400> 23

Ser Gly Ser Ser Ser Asn Ile Gly Ile Asn Ala Val Asn
1 5 10

<210> 24

<211> 6

<212> PRT

<213> Homo sapiens

<400> 24

Gly Asn Asp Gln Arg Pro
1 5

<210> 25

<211> 11

<212> PRT

<213> Homo sapiens

<400> 25

Ala Ala Trp Asp Asp Ser Leu Ser Gly Pro Val
1 5 10

<210> 26

<211> 17

<212> PRT

<213> Homo sapiens

<400> 26

Ala Ile Ser Gly Ser Gln Gly Ser Thr Tyr Tyr Ala Asp Ser Val Lys
1 5 10 15

Gly

<210> 27

<211> 17

<212> PRT

<213> Homo sapiens

<400> 27

Ala Ile Ser Gly Ser Gly Gly Ser Thr Trp Tyr Ala Asp Ser Val Lys
1 5 10 15

Gly

<210> 28

<211> 108

<212> PRT

<213> Homo sapiens

<400> 28

Leu Val Gln Pro Gly Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly
1 5 10 15

Phe Thr Phe Ser Ser Tyr Ala Met Ser Trp Val Arg Gln Ala Pro Gly

10

20 25 30

Lys Gly Leu Glu Trp Val Ser Ala Ile Ser Gly Ser Gly Gly Ser Thr
 35 40 45

Tyr Tyr Ala Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn
 50 55 60

Ser Lys Asn Thr Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp
 65 70 75 80

Thr Ala Val Tyr Tyr Cys Ala Arg Asp Asp Asp Tyr Asp Phe Asp
 85 90 95

Tyr Trp Gly Arg Gly Thr Met Val Thr Val Ser Ser
 100 105

<210> 29
 <211> 111
 <212> PRT
 <213> Homo sapiens

<400> 29
 Gln Ser Val Leu Thr Gln Pro Pro Ser Ala Ser Gly Thr Pro Gly Gln
 1 5 10 15

Arg Val Thr Ile Ser Cys Ser Gly Ser Ser Ser Asn Ile Gly Ile Asn
 20 25 30

Ala Val Asn Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu Leu
 35 40 45

Ile Tyr Gly Asn Asp Gln Arg Pro Ser Gly Val Pro Asp Arg Phe Ser
 50 55 60

Gly Ser Lys Ser Gly Thr Ser Ala Ser Leu Ala Ile Ser Gly Leu Gln
 65 70 75 80

Ser Glu Asp Glu Ala Asp Tyr Tyr Cys Ala Ala Trp Asp Asp Ser Leu
 85 90 95

Ser Gly Pro Val Phe Gly Gly Gly Thr Lys Leu Thr Val Leu Gly
 100 105 110

<210> 30
 <211> 235
 <212> PRT
 <213> Homo sapiens

<400> 30
 Leu Val Gln Pro Gly Gly Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly
 1 5 10 15

Phe Thr Phe Ser Ser Tyr Ala Met Ser Trp Val Arg Gln Ala Pro Gly
 20 25 30

Lys Gly Leu Glu Trp Val Ser Ala Ile Ser Gly Ser Gly Gly Ser Thr
 35 40 45

11

Tyr Tyr Ala Asp Ser Val Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn
 50 55 60
 Ser Lys Asn Thr Leu Tyr Leu Gln Met Asn Ser Leu Arg Ala Glu Asp
 65 70 75 80
 Thr Ala Val Tyr Tyr Cys Ala Arg Asp Asp Asp Tyr Asp Phe Asp
 85 90 95
 Tyr Trp Gly Arg Gly Thr Met Val Thr Val Ser Ser Gly Gly Gly Gly
 100 105 110
 Ser Gly Gly Gly Gly Ser Gly Gly Gly Gly Ser Ala Gln Ser Val Leu
 115 120 125
 Thr Gln Pro Pro Ser Ala Ser Gly Thr Pro Gly Gln Arg Val Thr Ile
 130 135 140
 Ser Cys Ser Gly Ser Ser Ser Asn Ile Gly Ile Asn Ala Val Asn Trp
 145 150 155 160
 Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu Leu Ile Tyr Gly Asn
 165 170 175
 Asp Gln Arg Pro Ser Gly Val Pro Asp Arg Phe Ser Gly Ser Lys Ser
 180 185 190
 Gly Thr Ser Ala Ser Leu Ala Ile Ser Gly Leu Gln Ser Glu Asp Glu
 195 200 205
 Ala Asp Tyr Tyr Cys Ala Ala Trp Asp Asp Ser Leu Ser Gly Pro Val
 210 215 220
 Phe Gly Gly Gly Thr Lys Leu Thr Val Leu Gly
 225 230 235

<210> 31
 <211> 14
 <212> PRT
 <213> Homo sapiens

<400> 31
 Tyr Phe Gly Lys Leu Glu Ser Lys Leu Ser Val Ile Arg Asn
 1 5 10

<210> 32
 <211> 14
 <212> PRT
 <213> Homo sapiens

<400> 32
 Glu Ser Lys Leu Ser Val Ile Arg Asn Leu Asn Asp Gln Val
 1 5 10

<210> 33
 <211> 14
 <212> PRT
 <213> Homo sapiens

<400> 33

Val Ile Arg Asn Leu Asn Asp Gln Val Leu Phe Ile Asp Gln
1 5 10

<210> 34

<211> 14

<212> PRT

<213> Homo sapiens

<400> 34

Asn Asp Gln Val Leu Phe Ile Asp Gln Gly Asn Arg Pro Leu
1 5 10

<210> 35

<211> 14

<212> PRT

<213> Homo sapiens

<400> 35

Phe Ile Asp Gln Gly Asn Arg Pro Leu Phe Glu Asp Met Thr
1 5 10

<210> 36

<211> 14

<212> PRT

<213> Homo sapiens

<400> 36

Asn Arg Pro Leu Phe Glu Asp Met Thr Asp Ser Asp Cys Arg
1 5 10

<210> 37

<211> 14

<212> PRT

<213> Homo sapiens

<400> 37

Glu Asp Met Thr Asp Ser Asp Cys Arg Asp Asn Ala Pro Arg
1 5 10

<210> 38

<211> 14

<212> PRT

<213> Homo sapiens

<400> 38

Ser Asp Cys Arg Asp Asn Ala Pro Arg Thr Ile Phe Ile Ile
1 5 10

<210> 39

<211> 14

<212> PRT

<213> Homo sapiens

13

<400> 39

Asn Ala Pro Arg Thr Ile Phe Ile Ile Ser Met Tyr Lys Asp
1 5 10

<210> 40

<211> 14

<212> PRT

<213> Homo sapiens

<400> 40

Ile Phe Ile Ile Ser Met Tyr Lys Asp Ser Gln Pro Arg Gly
1 5 10

<210> 41

<211> 14

<212> PRT

<213> Homo sapiens

<400> 41

Met Tyr Lys Asp Ser Gln Pro Arg Gly Met Ala Val Thr Ile
1 5 10

<210> 42

<211> 14

<212> PRT

<213> Homo sapiens

<400> 42

Gln Pro Arg Gly Met Ala Val Thr Ile Ser Val Lys Cys Glu
1 5 10

<210> 43

<211> 14

<212> PRT

<213> Homo sapiens

<400> 43

Ala Val Thr Ile Ser Val Lys Cys Glu Lys Ile Ser Thr Leu
1 5 10

<210> 44

<211> 14

<212> PRT

<213> Homo sapiens

<400> 44

Val Lys Cys Glu Lys Ile Ser Thr Leu Ser Cys Glu Asn Lys
1 5 10

<210> 45

<211> 14

<212> PRT

<213> Homo sapiens

<400> 45

14

Ile Ser Thr Leu Ser Cys Glu Asn Lys Ile Ile Ser Phe Lys
1 5 10

<210> 46

<211> 14

<212> PRT

<213> Homo sapiens

<400> 46

Cys Glu Asn Lys Ile Ile Ser Phe Lys Glu Met Asn Pro Pro
1 5 10

<210> 47

<211> 14

<212> PRT

<213> Homo sapiens

<400> 47

Ile Ser Phe Lys Glu Met Asn Pro Pro Asp Asn Ile Lys Asp
1 5 10

<210> 48

<211> 14

<212> PRT

<213> Homo sapiens

<400> 48

Met Asn Pro Pro Asp Asn Ile Lys Asp Thr Lys Ser Asp Ile
1 5 10

<210> 49

<211> 14

<212> PRT

<213> Homo sapiens

<400> 49

Asn Ile Lys Asp Thr Lys Ser Asp Ile Ile Phe Phe Gln Arg
1 5 10

<210> 50

<211> 14

<212> PRT

<213> Homo sapiens

<400> 50

Lys Ser Asp Ile Ile Phe Phe Gln Arg Ser Val Pro Gly His
1 5 10

<210> 51

<211> 14

<212> PRT

<213> Homo sapiens

<400> 51

Phe Phe Gln Arg Ser Val Pro Gly His Asp Asn Lys Met Gln
1 5 10

<210> 52
<211> 14
<212> PRT
<213> Homo sapiens

<400> 52
Val Pro Gly His Asp Asn Lys Met Gln Phe Glu Ser Ser Ser
1 5 10

<210> 53
<211> 14
<212> PRT
<213> Homo sapiens

<400> 54
Asn Lys Met Gln Phe Glu Ser Ser Ser Tyr Glu Gly Tyr Phe
1 5 10

<210> 54
<211> 14
<212> PRT
<213> Homo sapiens

<400> 54
Glu Ser Ser Ser Tyr Glu Gly Tyr Phe Leu Ala Cys Glu Lys
1 5 10

<210> 55
<211> 14
<212> PRT
<213> Homo sapiens

<400> 55
Glu Gly Tyr Phe Leu Ala Cys Glu Lys Glu Arg Asp Leu Phe
1 5 10

<210> 56
<211> 14
<212> PRT
<213> Homo sapiens

<400> 56
Ala Cys Glu Lys Glu Arg Asp Leu Phe Lys Leu Ile Leu Lys
1 5 10

<210> 57
<211> 14
<212> PRT
<213> Homo sapiens

<400> 57
Arg Asp Leu Phe Lys Leu Ile Leu Lys Lys Glu Asp Glu Leu
1 5 10

<210> 58
 <211> 14
 <212> PRT
 <213> Homo sapiens

<400> 58
 Leu Ile Leu Lys Lys Glu Asp Glu Leu Gly Asp Arg Ser Ile
 1 5 10

<210> 59
 <211> 14
 <212> PRT
 <213> Homo sapiens

<400> 59
 Glu Asp Glu Leu Gly Asp Arg Ser Ile Met Phe Thr Val Gln
 1 5 10

<210> 60
 <211> 12
 <212> PRT
 <213> Homo sapiens

<400> 60
 Asp Arg Ser Ile Met Phe Thr Val Gln Asn Glu Asp
 1 5 10

<210> 61
 <211> 157
 <212> PRT
 <213> Homo sapiens

<400> 61
 Tyr Phe Gly Lys Leu Glu Ser Lys Leu Ser Val Ile Arg Asn Leu Asn
 1 5 10 15

Asp Gln Val Leu Phe Ile Asp Gln Gly Asn Arg Pro Leu Phe Glu Asp
 20 25 30

Met Thr Asp Ser Asp Cys Arg Asp Asn Ala Pro Arg Thr Ile Phe Ile
 35 40 45

Ile Ser Met Tyr Lys Asp Ser Gln Pro Arg Gly Met Ala Val Thr Ile
 50 55 60

Ser Val Lys Cys Glu Lys Ile Ser Thr Leu Ser Cys Glu Asn Lys Ile
 65 70 75 80

Ile Ser Phe Lys Glu Met Asn Pro Pro Asp Asn Ile Lys Asp Thr Lys
 85 90 95

Ser Asp Ile Ile Phe Phe Gln Arg Ser Val Pro Gly His Asp Asn Lys
 100 105 110

Met Gln Phe Glu Ser Ser Ser Tyr Glu Gly Tyr Phe Leu Ala Cys Glu
 115 120 125

17

Lys Glu Arg Asp Leu Phe Lys Leu Ile Leu Lys Lys Glu Asp Glu Leu
 130 135 140

Gly Asp Arg Ser Ile Met Phe Thr Val Gln Asn Glu Asp
 145 150 155

<210> 62
 <211> 341
 <212> DNA
 <213> Homo sapiens

<220>
 <221> CDS
 <222> (1)..(339)

<400> 62
 cag gtc cag ctg gtg cag tct ggg gct gag gtg aag aag cct ggg gcc 48
 Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
 1 5 10 15
 tcg atg aaa gtc tcc tgt aag act tct gga tac acc ttc acc ggc tat 96
 Ser Met Lys Val Ser Cys Lys Thr Ser Gly Tyr Thr Phe Thr Gly Tyr
 20 25 30
 tat atc cac tgg gtg cga cag gcc cct gga cag gga ttc gag tgg ata 144
 Tyr Ile His Trp Val Arg Gln Ala Pro Gly Gln Gly Phe Glu Trp Ile
 35 40 45
 gga cgg ctc aac ccc acc act ggt gac gca aat ttt gca gaa aag ttt 192
 Gly Arg Leu Asn Pro Thr Thr Gly Asp Ala Asn Phe Ala Glu Lys Phe
 50 55 60
 cag ggc agg gtc gcc ctg acc aga gac acg tcc atc agc aca gcc tat 240
 Gln Gly Arg Val Ala Leu Thr Arg Asp Thr Ser Ile Ser Thr Ala Tyr
 65 70 75 80
 tta caa cta gac agc ctc aaa tct gac gac acg gcc gta tat tat tgt 288
 Leu Gln Leu Asp Ser Leu Lys Ser Asp Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
 gcg gga aaa gag ggt gcc tgg ggc cag ggc acc ctg gtc acc gtc tcg 336
 Ala Gly Lys Glu Gly Ala Trp Gly Gln Gly Thr Leu Val Thr Val Ser
 100 105 110
 agt gg 341
 Ser

<210> 63
 <211> 113
 <212> PRT
 <213> Homo sapiens

<400> 63
 Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala
 1 5 10 15
 Ser Met Lys Val Ser Cys Lys Thr Ser Gly Tyr Thr Phe Thr Gly Tyr
 20 25 30

18

Tyr Ile His Trp Val Arg Gln Ala Pro Gly Gln Gly Phe Glu Trp Ile
 35 40 45

Gly Arg Leu Asn Pro Thr Thr Gly Asp Ala Asn Phe Ala Glu Lys Phe
 50 55 60

Gln Gly Arg Val Ala Leu Thr Arg Asp Thr Ser Ile Ser Thr Ala Tyr
 65 70 75 80

Leu Gln Leu Asp Ser Leu Lys Ser Asp Asp Thr Ala Val Tyr Tyr Cys
 85 90 95

Ala Gly Lys Glu Gly Ala Trp Gly Gln Gly Thr Leu Val Thr Val Ser
 100 105 110

Ser

<210> 64
 <211> 327
 <212> DNA
 <213> Homo sapiens

<220>
 <221> CDS
 <222> (1)..(327)

<400> 64
 tct tct gag ctg act cag gac cct gct gtg tct gtg gcc ttg gga cag 48
 Ser Ser Glu Leu Thr Gln Asp Pro Ala Val Ser Val Ala Leu Gly Gln
 1 5 10 15

aca gtc agg atc aca tgc caa gga gac agc ctc aga cac ttt tat cca 96
 Thr Val Arg Ile Thr Cys Gln Gly Asp Ser Leu Arg His Phe Tyr Pro
 20 25 30

aac tgg tac cag cag aag cca gga cag gcc cct gta ctt gtc atc tat 144
 Asn Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Val Leu Val Ile Tyr
 35 40 45

ggt aaa aac aat cgg ccc tca ggg atc cca gac cga ttc tct ggc tcc 192
 Gly Lys Asn Asn Arg Pro Ser Gly Ile Pro Asp Arg Phe Ser Gly Ser
 50 55 60

ggc tca gga aac aca ggt tcc ttg acc atc act ggg gcc cag gcg gaa 240
 Gly Ser Gly Asn Thr Gly Ser Leu Thr Ile Thr Gly Ala Gln Ala Glu
 65 70 75 80

gat gag gct gac tat tac tgt ggc tcc cgg gac agc agt ggt atc cat 288
 Asp Glu Ala Asp Tyr Tyr Cys Gly Ser Arg Asp Ser Ser Gly Ile His
 85 90 95

gtg gta ttc ggc gga ggg acc aag gtc acc gtc cta ggt 327
 Val Val Phe Gly Gly Gly Thr Lys Val Thr Val Leu Gly
 100 105

<210> 65
 <211> 109
 <212> PRT

<213> Homo sapiens

<400> 65

```

Ser Ser Glu Leu Thr Gln Asp Pro Ala Val Ser Val Ala Leu Gly Gln
 1             5             10             15

Thr Val Arg Ile Thr Cys Gln Gly Asp Ser Leu Arg His Phe Tyr Pro
      20             25             30

Asn Trp Tyr Gln Gln Lys Pro Gly Gln Ala Pro Val Leu Val Ile Tyr
      35             40             45

Gly Lys Asn Asn Arg Pro Ser Gly Ile Pro Asp Arg Phe Ser Gly Ser
 50             55             60

Gly Ser Gly Asn Thr Gly Ser Leu Thr Ile Thr Gly Ala Gln Ala Glu
 65             70             75             80

Asp Glu Ala Asp Tyr Tyr Cys Gly Ser Arg Asp Ser Ser Gly Ile His
      85             90             95

Val Val Phe Gly Gly Gly Thr Lys Val Thr Val Leu Gly
      100             105

```

<210> 66

<211> 354

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (1)..(354)

<400> 66

```

gag gtg cag ctg ttg gag tct ggg gga ggc ttg gta cag cct ggg ggg      48
Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
 1             5             10             15

tcc ctg aga ctc tcc tgt gca gcc tct gga ttc acc ttt agc agc tat      96
Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
      20             25             30

gcc atg agc tgg gtc cgc cag gct cca ggg aag ggg ctg gag tgg gtc      144
Ala Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
      35             40             45

tca gct att agt ggt agt ggt ggt agc aca tac tac gca gac tcc gtg      192
Ser Ala Ile Ser Gly Ser Gly Gly Ser Thr Tyr Tyr Ala Asp Ser Val
      50             55             60

aag ggc cgg ttc acc atc tcc aga gac aat tcc aag aac acg ctg tat      240
Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
      65             70             75             80

ctg caa atg aac agc ctg aga gcc gag gac acg gcc gtg tat tac tgt      288
Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
      85             90             95

gcg aga gat gac gat gac tac gac ttt gac tac tgg ggc cgg ggg aca      336
Ala Arg Asp Asp Asp Asp Tyr Asp Phe Asp Tyr Trp Gly Arg Gly Thr

```


100 105 20 110
 atg gtc acc gtc tcg agt 354
 Met Val Thr Val Ser Ser
 115
 <210> 67
 <211> 118
 <212> PRT
 <213> Homo sapiens
 <400> 67
 Glu Val Gln Leu Leu Glu Ser Gly Gly Gly Leu Val Gln Pro Gly Gly
 1 5 10 15
 Ser Leu Arg Leu Ser Cys Ala Ala Ser Gly Phe Thr Phe Ser Ser Tyr
 20 25 30
 Ala Met Ser Trp Val Arg Gln Ala Pro Gly Lys Gly Leu Glu Trp Val
 35 40 45
 Ser Ala Ile Ser Gly Ser Gly Gly Ser Thr Tyr Tyr Ala Asp Ser Val
 50 55 60
 Lys Gly Arg Phe Thr Ile Ser Arg Asp Asn Ser Lys Asn Thr Leu Tyr
 65 70 75 80
 Leu Gln Met Asn Ser Leu Arg Ala Glu Asp Thr Ala Val Tyr Tyr Cys
 85 90 95
 Ala Arg Asp Asp Asp Asp Tyr Asp Phe Asp Tyr Trp Gly Arg Gly Thr
 100 105 110
 Met Val Thr Val Ser Ser
 115
 <210> 68
 <211> 334
 <212> DNA
 <213> Homo sapiens
 <220>
 <221> CDS
 <222> (1)..(333)
 <400> 68
 cag tct gtg ttg acg cag ccg ccc tca gcg tct ggg gcc ccc ggt cag 48
 Gln Ser Val Leu Thr Gln Pro Pro Ser Ala Ser Gly Ala Pro Gly Gln
 1 5 10 15
 agg gtc acc atc tct tgt tct gga agc agc tcc aac atc gga att aat 96
 Arg Val Thr Ile Ser Cys Ser Gly Ser Ser Ser Asn Ile Gly Ile Asn
 20 25 30
 gct gta aac tgg tac cag cag ctc cca gga acg gcc ccc aaa ctc ctc 144
 Ala Val Asn Trp Tyr Gln Gln Leu Pro Gly Thr Ala Pro Lys Leu Leu
 35 40 45
 atc tat ggt aat gat cag cgg ccc tca ggg gtc cct gac cga ttc tct 192
 Ile Tyr Gly Asn Asp Gln Arg Pro Ser Gly Val Pro Asp Arg Phe Ser

```

<400> 70
Tyr Phe Gly Lys Leu Glu Ser Lys Leu Ser Val Ile Arg Asn Leu Asn
  1             5             10             15
Asp Gln Val Leu Phe Ile Asp Gln Gly Asn Arg Pro Leu Phe Glu Asp
      20             25             30
Met Thr Asp Ser Asp Cys Arg Asp Asn Ala Pro Arg Thr Ile Phe Ile
      35             40             45
Ile Ser Met Tyr Lys Asp Ser Gln Pro Arg Gly Met Ala Val Thr Ile
  50             55             60

```

Ser Val
65

<210> 71
<211> 34
<212> PRT
<213> Homo sapiens

<400> 71

Phe Leu Ala Cys Glu Lys Glu Arg Asp Leu Phe Lys Leu Ile Leu Lys
1 5 10 15

Lys Glu Asp Glu Leu Gly Asp Arg Ser Ile Met Phe Thr Val Gln Asn
20 25 30

Glu Asp

THIS PAGE BLANK (USPTO)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
16 August 2001 (16.08.2001)

PCT

(10) International Publication Number
WO 01/58956 A3

(51) International Patent Classification⁷: **C07K 16/24**,
A61K 39/395, C12N 5/10, 15/13, A61K 38/17, 38/13,
31/505, 31/445, C12N 15/85, A61P 1/00, 3/00, 5/00, 7/00,
9/00, 11/00, 13/00, 15/00, 17/00, 19/00, 21/00, 25/00,
29/00, 31/00, 33/00, 35/00, 37/00, A61K 31/56, 31/56,
31/56 // (A61K 39/395, 31:505) (A61K 39/395, 38:13)
(A61K 39/395, 31:445) (A61K 39/395, 31:56) (A61K
39/395, 38:17)

(21) International Application Number: PCT/US01/04170

(22) International Filing Date: 9 February 2001 (09.02.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/181,608 10 February 2000 (10.02.2000) US

(63) Related by continuation (CON) or continuation-in-part
(CIP) to earlier application:
US 60/181,608 (CON)
Filed on 10 February 2000 (10.02.2000)

(71) Applicant (for all designated States except US): **BASF
AKTIENGESELLSCHAFT** [DE/DE]; Ludwigshafen,
Rheinland-Pfalz (DE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **GHAYUR, Tariq**
[PK/US]; 1014 Washington Street, Holliston, MA
01746 (US). **DIXON, Richard, W.** [US/US]; 6 Samuel
Drive, North Grafton, MA 01546 (US). **ROGUSKA,**
Mike [US/US]; 16 Hilldale Road, Ashland, MA 01721
(US). **WHITE, Michael** [US/US]; 30 Angelica Drive,
Framingham, MA 01701 (US). **LABKOVSKY, Boris**
[US/US]; 107A-5 Broadmeadow Road, Marlborough,
MA 01752 (US). **SALFELD, Jochen** [DE/US]; 177 Old

Westboro Road, North Grafton, MA 01536 (US). **DUN-
CAN, Alexander, Robert** [GB/GB]; 8 Garden Fields,
Little Shelford, Cambridge CB2 5HH (GB). **BROCK-
LEHURST, Simon, Mark** [GB/GB]; 38 Huntmill,
Fulbourn, Cambridge CB1 5RH (GB). **MANKOVICH,**
John [US/US]; 416 Lowell Street, Andover, MA 01810
(US). **SHORROCK, Celia, Patricia** [GB/US]; 17 Stanley
Road, Cambridge CB5 8LF (GB). **THOMPSON, Julia,**
Elizabeth [GB/GB]; Seeby's Cottage, 6 High Street,
Whittlesford, Cambridge CB2 4LT (GB). **LENNARD,**
Simon, Nicholas [GB/GB]; 1 Bakers Lane, Linton, Cam-
bridgeshire CB1 6NF (GB).

(74) Agents: **DECONTI, Giulio, A., JR.** et al.; Lahive &
Cockfield, LLP, 28 State Street, Boston, MA 02109 (US).

(81) Designated States (national): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,
TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

(88) Date of publication of the international search report:
7 March 2002

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: ANTIBODIES THAT BIND HUMAN INTERLEUKIN-18 AND METHODS OF MAKING AND USING

(57) Abstract: Antibodies that bind human interleukin-18 (hIL-18) are provided, in particular antibodies that bind epitope(s) of human IL-18. The antibodies can be, for example, entirely human antibodies, recombinant antibodies, or monoclonal antibodies. Preferred antibodies have high affinity for hIL-18 and neutralize hIL-18 activity *in vitro* and *in vivo*. An antibody of the invention can be a full-length antibody or an antigen-binding portion thereof. Method of making and method of using the antibodies of the invention are also provided. The antibodies, or antibody portions, of the invention are useful for detecting hIL-18 and for inhibiting hIL-18 activity, e.g., in a human subject suffering from a disorder in which hIL-18 activity is detrimental.

WO 01/58956 A3

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 01/04170

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7	C07K16/24	A61K39/395	C12N5/10	C12N15/13	A61K38/17
	A61K38/13	A61K31/505	A61K31/445	C12N15/85	A61P1/00
	A61P3/00	A61P5/00	A61P7/00	A61P9/00	A61P11/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07K A61K C12N A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, BIOSIS, WPI Data, PAJ

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 974 600 A (HAYASHIBARA BIOCHEM LAB) 26 January 2000 (2000-01-26) paragraph [0021] paragraph [0038] - paragraph [0040] paragraph [0045] - paragraph [0071] claims 1-27	1-3,53, 55,57,59
X	KOHKA HIDEO ET AL: "Involvement of interleukin-18 (IL-18) in mixed lymphocyte reactions (MLR)." JOURNAL OF INTERFERON AND CYTOKINE RESEARCH, vol. 19, no. 9, 1999, pages 1053-1057, XP001007317 ISSN: 1079-9907 the whole document	1,2,53, 55,57, 59,60



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

24 July 2001

Date of mailing of the international search report

17.09.01

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl.
Fax: (+31-70) 340-3016

Authorized officer

Renggli, J

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 01/04170

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61P13/00 A61P15/00 A61P17/00 A61P19/00 A61P21/00
A61P25/00 A61P29/00 A61P31/00 A61P33/00 A61P35/00
A61P37/00 A61K31/56 A61K31/56 A61K31/56

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category * Citation of document, with indication, where appropriate, of the relevant passages Relevant to claim No.

P,X	WO 00 56771 A (HOLMES STEPHEN D ;SMITHKLINE BEECHAM PLC (GB); ABDEL MEGUID SHERIN) 28 September 2000 (2000-09-28) page 2 -page 3; claims 1-23 -----	1,2,53, 55,57,59
-----	---	---------------------

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance
"E" earlier document but published on or after the international filing date
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
"O" document referring to an oral disclosure, use, exhibition or other means
"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
"G" document member of the same patent family

Date of the actual completion of the international search

24 July 2001

Date of mailing of the international search report

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl.
Fax: (+31-70) 340-3016

Authorized officer

Renggli, J

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 01/04170

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 // (A61K39/395,31:505), (A61K39/395,38:13), (A61K39/395,31:445),
(A61K39/395,31:56), (A61K39/395,38:17)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"Z" document member of the same patent family

Date of the actual completion of the international search

24 July 2001

Date of mailing of the international search report

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Renggli, J

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US 01/04170

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Although claims 53, 55, 57, 59 and 60 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

1-3, 53, 55, 57, 59 (part) and 60

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 1-3, 53, 55, 57, 59 (part) and 60

A compound capable of binding a human IL-18 amino acid sequence or portion thereof, wherein said amino acid comprises a sequence selected for the group consisting of SEQ ID NO:67 and SEQ ID NO:68 and methods of inhibiting or treating involving the use of said compound.

2. Claims: 4-10, 39-46 (part), 54 (part), 56 (part), 58 (part) and 59 (part)

A human monoclonal antibody, or antigen binding portion thereof, capable of binding to human IL-18 and isolated nucleic acid encoding an CDR of said antibody, host cells containing the said nucleic acid, method of synthesizing said antibody, pharmaceutical compositions comprising said antibody, methods for inhibiting or treating involving the use of said antibody.

3. Claims: 11-15, 39-46 (part), 54 (part), 56 (part), 58 (part) and 59 (part)

An isolated antibody, or an antigen-binding portion thereof, that binds an epitope of human IL-18, or portion thereof, comprising an amino acid sequence selected from the group comprising seq ID:3 and 33 and isolated nucleic acid encoding an CDR of said antibody, host cells containing the said nucleic acid, method of synthesizing said antibody, pharmaceutical compositions comprising said antibody, methods for inhibiting or treating involving the use of said antibody.

4. Claims: 16-21, 39-46 (part), 54 (part), 56 (part), 58 (part) and 59 (part)

An isolated antibody, or antigen-binding portion thereof, that binds to an epitope of human IL-18, wherein the antibody, or antigen-binding portion thereof, dissociates from human IL-18 with a koff rate constant of 0.1 s⁻¹ or less, as determined by surface plasmon resonance, or which inhibits human IL-18 activity with an IC₅₀ of 1 x 10⁻⁶ M or less and isolated nucleic acid encoding an CDR of said antibody, host cells containing the said nucleic acid, method of synthesizing said antibody, pharmaceutical compositions comprising said antibody, methods for inhibiting or treating involving the use of said antibody.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

5. Claims: 22-28, 32-35, 39-46 (part), 54 (part), 56 (part),
58 (part) and 59 (part)

An isolated human antibody, or an antigen-binding portion thereof, comprising at least one variable region CDR domain capable of binding an epitope of human IL-18 and isolated nucleic acid encoding an CDR of said antibody, host cells containing the said nucleic acid, method of synthesizing said antibody, pharmaceutical compositions comprising said antibody, methods for inhibiting or treating involving the use of said antibody.

6. Claims: 29-31, 39-46 (part), 54 (part), 56 (part),
58 (part) and 59 (part)

An isolated antibody, or an antigen-binding portion thereof, with a variable region comprising an amino acid sequence selected from the group consisting of seq IDs: 15, 16 and 17 and isolated nucleic acid encoding an CDR of said antibody, host cells containing the said nucleic acid, method of synthesizing said antibody, pharmaceutical compositions comprising said antibody, methods for inhibiting or treating involving the use of said antibody.

7. Claims: 36-38, 39-46 (part), 54 (part), 56 (part),
58 (part) and 59 (part)

An isolated antibody, or an antigen-binding portion thereof, with a variable region comprising an amino acid selected from the group consisting of seq IDs: 26, 27 and 29 and isolated nucleic acid encoding an CDR of said antibody, host cells containing the said nucleic acid, method of synthesizing said antibody, pharmaceutical compositions comprising said antibody, methods for inhibiting or treating involving the use of said antibody.

8. Claims: 47-52

A method of making an antibody that binds human interleukin-18 comprising exposing an antibody repertoire to an antigen comprising an epitope of human IL-18 or portion thereof and selecting from the antibody repertoire an antibody that binds the epitope of human IL-18, or portion thereof.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 01/04170

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 0974600	A	26-01-2000	JP 2000236884 A	05-09-2000
WO 0056771	A	28-09-2000	NONE	

Form PCT/ISA/210 (patent family annex) (July 1992)

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☒ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

THIS PAGE BLANK (US)